

# *Pseudo-nitzschia* species (Bacillariophyceae) identification and delineation using mitochondrial *cox1* gene sequences as compared to LSU rDNA

Suh Nih Tan<sup>1</sup> · Hong Chang Lim<sup>2</sup> · Sing Tung Teng<sup>3</sup> · Po Teen Lim<sup>1</sup> · Chui Pin Leaw<sup>1</sup>

Received: 11 February 2015 / Accepted: 9 June 2015 / Published online: 28 July 2015  
© Japanese Society of Fisheries Science 2015

**Abstract** An attempt was made to infer the phylogeny of *Pseudo-nitzschia* species by using the mitochondrial-encoded gene, cytochrome c oxidase subunit I (*cox1*), and comparing it with the nuclear-encoded large subunit ribosomal DNA (LSU rDNA). A pair of primers targeting *Pseudo-nitzschia cox1* was designed in silico and used to infer the molecular phylogeny of *Pseudo-nitzschia*. The primer pair was tested using genomic DNAs isolated from six species of *Pseudo-nitzschia* from Malaysia. The phylogenetic inference of *cox1* was then compared to the LSU rDNA phylogeny. Phylogenetic reconstructions of both data sets revealed monophyly of *Pseudo-nitzschia* species complexes. The range of genetic divergences among *Pseudo-nitzschia* species were higher in the *cox1* data set (3.5–20.4 %) compared to the LSU rDNA data set (0.1–8.8 %). The present study suggests that high genetic divergence in *cox1* of *Pseudo-nitzschia* species could be a useful genetic marker for DNA bar coding.

**Keywords** *cox1* · Diatom · LSU rDNA · Primer · *Pseudo-nitzschia*

## Introduction

Diatoms are an important component of the marine phytoplankton, in terms of both abundance and species richness [1]. While most diatom species are important ecologically, several species are known to cause amnesic shellfish poisoning (ASP). The first incident of ASP was reported in Canada, in 1987; three deaths and 105 cases of acute human intoxication were documented after consuming the contaminated blue mussels (*Mytilus edulis* Linnaeus) [2]. The neurotoxin, domoic acid (DA), C<sub>15</sub>H<sub>21</sub>NO<sub>6</sub>, was later found to be produced by the diatom, *Pseudo-nitzschia multiseriis* (Hasle) Hasle (formerly known as *Nitzschia pungens* f. *multiseriis*) [3]. People who consumed the contaminated shellfish in relation to ASP showed similar symptoms to Alzheimer's disease, i.e. loss of short-term memory [2]; other symptoms included diarrhea, nausea, vomiting, abdominal cramps and headache.

In Malaysia, there have been no reported cases of ASP. Screening of ASP-toxin production among the cultured strains of *P. pungens* (Grunow ex Cleve) Hasle showed undetectable levels of DA [4], even though the species is commonly found in Malaysian waters [5, 6]. High species diversity of *Pseudo-nitzschia* species in Malaysian coastal waters was reported by Teng et al. [5], with more than 23 species documented. Among the 23 species, six species have been identified from the waters as new morphotypes, and described as *P. batesiana* H.C. Lim, Teng, Leaw et P.T. Lim, *P. circumspora* H.C. Lim, Leaw et P.T. Lim, *P. fukuyoi* H.C. Lim, Teng, Leaw et P.T. Lim, *P. kodamae* Teng, H.C. Lim, Leaw et P.T. Lim, *P. lundholmiae* H.C. Lim, Teng,

**Electronic supplementary material** The online version of this article (doi:10.1007/s12562-015-0902-7) contains supplementary material, which is available to authorized users.

✉ Chui Pin Leaw  
cpleaw@um.edu.my; chuiplinleaw@gmail.com

<sup>1</sup> Bachok Marine Research Station, Institute of Ocean and Earth Sciences, University of Malaya, 16310 Bachok, Kelantan, Malaysia

<sup>2</sup> Tunku Abdul Rahman University College, Johor Branch Campus, 85000 Segamat, Johor, Malaysia

<sup>3</sup> Faculty of Resource Science and Technology, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia