

Lost Aroids: On the taxonomic importance of relocating poorly collected species

PETER C. BOYCE¹ and WONG SIN YENG²

Aridarum montanum Ridl. and *Piptospatha insignis* N.E.Br. (Araceae: Schismatoglottideae), aroids originating from Borneo that are each known from a single collection, are discussed and illustrated. The history of their discovery is reviewed, together with what is known or speculated of their ecology. The biological significance of the collection locality of *A. montanum* is highlighted. The species' individual importance to modern systematics is emphasized.

Keywords. Araceae, *Aridarum*, *Piptospatha*, Borneo, Malaysia, Sarawak, Santubong

INTRODUCTION

'Lost' plant species – species tantalizingly only known from a single herbarium collection, or frustratingly from just an old illustration, hold an abiding fascination for plant enthusiasts, whether professional botanists or keen hobbyists. The ranks of these 'still lost' plant species are perhaps no better exemplified than by *Archivea kewensis* Christenson & Jenny (Fig. 1), a Brazilian species (indeed, a genus) known from a single 19th century watercolour deposited in the Herbarium & Archives of Kew Gardens, from whence the genus and species epithets are derived (Christenson & Jenny, 1996).

Aroids, perhaps by reason of their often originating from almost inaccessible tropical forests, are host to a remarkable number of such 'lost' species. Remarkable, too, is that quite some number of long-lost species has been re-found over the past 20 years. Of particular note [with the period "lost" in years] are: *Gearum brasiliense* N.E.Br. [150 years] (Mayo et al., 1994), *Mangonia tweediana* Schott [142 years] (Bogner & Marchesi, 2000), *Zomicarpella maculata* N.E.Br. [116 years] (Bogner, 2007, 2009), and *Ulearum sagittatum* Engl. [90 years] (Boyce, 1995; Bogner, 1997).

¹ Manuscript submitted: 9 November 2011

Manuscript accepted: 23 November 2011

Pusat Pengajian Sains Kajihayat, Universiti Sains Malaysia 11800 USM, Pulau Pinang, Malaysia

² Department of Plant Science & Environmental Ecology, Faculty of Resource Science & Technology, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia

* Corresponding author email: phymatarum@gmail.com