Clinical Microbiology Reviews

Human Infections and Detection of Plasmodium knowlesi

Balbir Singh and Cyrus Daneshvar *Clin. Microbiol. Rev.* 2013, 26(2):165. DOI: 10.1128/CMR.00079-12.

Updated information and services can be found at: http://cmr.asm.org/content/26/2/165
These include:
This article cites 134 articles, 33 of which can be accessed free at: http://cmr.asm.org/content/26/2/165#ref-list-1

CONTENT ALERTS

REFERENCES

Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), more»

Information about commercial reprint orders: http://journals.asm.org/site/misc/reprints.xhtml To subscribe to to another ASM Journal go to: http://journals.asm.org/site/subscriptions/



Human Infections and Detection of Plasmodium knowlesi

Balbir Singh, a Cyrus Daneshvarb

Malaria Research Centre, Universiti Malaysia Sarawak, Kuching, Sarawak, Malaysia^a; Oxford Centre for Respiratory Medicine, Oxford University Hospitals, Headington, United Kingdomb

SUMMARY	165
INTRODUCTION	165
Life Cycle of <i>Plasmodium</i> in Humans	166
Discovery and Early Studies	
EPIDEMIOLOGY	
Distribution	
Reservoir Hosts	
Vectors	
Populations at Risk	
Molecular Epidemiology and Evolutionary and Demographic History	
LABORATORY DIAGNOSIS	
Microscopy	169
Molecular Detection Methods.	
Rapid Diagnostic Tests	172
CLINICAL COURSE	173
Historical Data	173
Recent Studies.	174
Demographics	174
Symptoms	174
Clinical Examination Findings	174
Laboratory Findings	
Thrombocytopenia	174
Anemia	
Other hematological findings	
Renal function	
Liver function	
Clinical Aspects in Children	
Complicated Knowlesi Malaria	
Severe disease	
Acute respiratory distress syndrome.	
Acute renal failure	
Other features	
Pathogenesis	
TREATMENT	
FUTURE DIRECTIONS AND CHALLENGES	
ACKNOWLEDGMENTS	
REFERENCES	
AUTHOR BIOS	184

SUMMARY

Plasmodium knowlesi is a malaria parasite that is found in nature in long-tailed and pig-tailed macaques. Naturally acquired human infections were thought to be extremely rare until a large focus of human infections was reported in 2004 in Sarawak, Malaysian Borneo. Human infections have since been described throughout Southeast Asia, and P. knowlesi is now recognized as the fifth species of Plasmodium causing malaria in humans. The molecular, entomological, and epidemiological data indicate that human infections with P. knowlesi are not newly emergent and that knowlesi malaria is primarily a zoonosis. Human infections were undiagnosed until molecular detection methods that could distinguish P. knowlesi from the morphologically similar human malaria parasite P. malariae became available. P. knowlesi infections cause a spectrum of disease and are potentially fatal, but if detected early enough, infections in humans are readily treatable. In this review on knowlesi malaria, we describe the early studies on P. knowlesi and focus on the epidemiology, diagnosis, clinical aspects, and treatment of knowlesi malaria. We also discuss the gaps in our knowledge and the challenges that lie ahead in studying the epidemiology and pathogenesis of knowlesi malaria and in the prevention and control of this zoonotic infection.

INTRODUCTION

alaria is caused by protozoan parasites belonging to the genus Plasmodium. Over 150 species have been described to date, infecting mammals, birds, and reptiles (1). Despite having

Address correspondence to Balbir Singh, bskhaira55@gmail.com. Copyright © 2013, American Society for Microbiology. All Rights Reserved. doi:10.1128/CMR.00079-12