



Faculty of Resource Science and Technology

Isolation and Characterization of Methicillin-Resistant *Staphylococcus aureus* from Air Conditioning System in Faculty of Resource Science and Technology, Universiti Malaysia Sarawak

Nurul Nadia Binti Zayadi

(53489)

**Bachelor of Science with Honours
(Resource Biotechnology)
2018**

**Isolation and Characterization of Methicillin-Resistant *Staphylococcus aureus*
from Air Conditioning System in Faculty of Resource Science and Technology,
Universiti Malaysia Sarawak**

Nurul Nadia Binti Zayadi (53489)

A thesis submitted in fulfilment of the Final Year Project (STF 3013) Course

Faculty of Resource Science and Technology

UNIVERSITI MALAYSIA SARAWAK

2018

UNIVERSITI MALAYSIA SARAWAK

Grade: _____

Please tick (✓)

Final Year Project Report

Masters

PhD

DECLARATION OF ORIGINAL WORK

This declaration is made on the 11 day of June year 2018

Student's Declaration:

I Nurul Nadia Bt. Dayadi (53489) (FSTS)

(PLEASE INDICATE NAME, MATRIC NO. AND FACULTY) hereby declare that the work entitled, Isolation and characterization of MRSA from air conditioning system in FST, UNIMAS. is my original work. I have not copied from any other students' work or from any other sources with the exception where due reference or acknowledgement is made explicitly in the text, nor has any part of the work been written for me by another person.

11/June/2018

Date submitted

Nurul Nadia Bt. Dayadi (53489)

Name of the student (Matric No.)

Supervisor's Declaration:

I, (SUPERVISOR'S NAME), hereby certify that the work entitled, (TITLE) was prepared by the aforementioned or above mentioned student, and was submitted to the "FACULTY" as a * partial/full fulfillment for the conferment of (PLEASE INDICATE THE DEGREE TITLE), and the aforementioned work, to the best of my knowledge, is the said student's work

Received for examination by: Prof. Dr. Kasing Apun Date: 11/6/2018 (Name of the supervisor)

I declare this Project/Thesis is classified as (Please tick (√)):

- CONFIDENTIAL** (Contains confidential information under the Official Secret Act 1972)*
 RESTRICTED (Contains restricted information as specified by the organisation where research was done)*
 OPEN ACCESS

I declare this Project/Thesis is to be submitted to the Centre for Academic Information Services (CAIS) and uploaded into UNIMAS Institutional Repository (UNIMAS IR) (Please tick (√)):


- YES**
 NO

Validation of Project/Thesis

I hereby duly affirmed with free consent and willingness declared that this said Project/Thesis shall be placed officially in the Centre for Academic Information Services with the abide interest and rights as follows:

- This Project/Thesis is the sole legal property of Universiti Malaysia Sarawak (UNIMAS).
- The Centre for Academic Information Services has the lawful right to make copies of the Project/Thesis for academic and research purposes only and not for other purposes.
- The Centre for Academic Information Services has the lawful right to digitize the content to be uploaded into Local Content Database.
- The Centre for Academic Information Services has the lawful right to make copies of the Project/Thesis if required for use by other parties for academic purposes or by other Higher Learning Institutes.
- No dispute or any claim shall arise from the student himself / herself neither a third party on this Project/Thesis once it becomes the sole property of UNIMAS.
- This Project/Thesis or any material, data and information related to it shall not be distributed, published or disclosed to any party by the student himself/herself without first obtaining approval from UNIMAS.

Student's signature 
91/7492/2018
(Date)

Supervisor's signature: 
(Date) 11/6/2018

Current Address:

Notes: * If the Project/Thesis is **CONFIDENTIAL** or **RESTRICTED**, please attach together as annexure a letter from the organisation with the date of restriction indicated, and the reasons for the confidentiality and restriction.

[The instrument was prepared by The Centre for Academic Information Services]

ACKNOWLEDGEMENTS

In the name of Allah, the Most Gracious and Most Merciful

First and foremost, I would like to give my greatest gratitude to Allah S.W.T for his blessing and grace for me in order to complete this thesis with courage and strength.

I would also like to take this chance to express my gratitude to my supervisor Professor Kasing Apun for her kind advices and numerous supports in order for me complete this thesis successfully. Without her guidance, it is impossible for me to complete this thesis all by myself. I am so grateful for having this chance to complete my Final Year Project under her supervision.

I am also very thankful to my lab mates, especially Laurie, Scholastica, Stephanie, and also Syazira. Their help for me in completing my thesis is a big help to me. They always give their support when I am having difficulties while doing my project. Furthermore, I am also very thankful to one of the master student who is Jennifer Jalan for assisting me throughout my project, and also for Mr Azis, the staff of FRST. Lastly, special thanks to my families and friends for their moral support, advises and financial support throughout the duration of my studies.

Isolation and Characterization of Methicillin-Resistant *Staphylococcus aureus* (MRSA)
from Air Conditioning System in Faculty of Resource Science and Technology,

Universiti Malaysia Sarawak

Nurul Nadia Binti Zayadi

Resource Biotechnology Programme
Faculty of Science and Technology
Universiti Malaysia Sarawak

ABSTRACT

Staphylococcus aureus (*S. aureus*) is one of major human pathogen that able cause mild to serious infections. Present treatment becomes challenging to control due to the presence of methicillin-resistant *Staphylococcus aureus* (MRSA). The present study looks on the isolation and characterization of MRSA from air conditioning system in Faculty of Resources Science and Technology, UNIMAS. The swab samples were isolated from the ventilation of central air conditioner and dipped into 0.1 % peptone water. The swab samples were then carried out for serial dilution and plating on mannitol salt agar (MSA) and incubated for 24-36 hours in room temperature. Suspected colonies of *S. aureus* appeared as whitish colonies on MSA agar, and have gram-positive cocci in clusters. Only 16.67% of the total samples with dilution factor of 10^{-5} were positive for suspected *S. aureus* and were chosen to proceed for the following biochemical test. However, only 25% of the isolates were tested positive for catalase test, gelatin liquefaction test, and methyl red test, except for citrate test and urease test. All of the isolates were then carried out to assess their resistance towards four antibiotics by using disc diffusion method. All of the isolates were considered as resistant towards methicillin, penicillin, vancomycin, and cephalothin. This study is significant to increase awareness to UNIMAS staffs, lecturers, and students, by providing basic data for UNIMAS authorities on the status of MRSA from air conditioning system in UNIMAS.

Key words: Methicillin-resistant *Staphylococcus aureus* (MRSA), mannitol salt agar (MSA), disc diffusion method.

ABSTRAK

Staphylococcus aureus ialah salah satu daripada patogen yang mampu menyebabkan infeksi yang sederhana kepada yang lebih serius. Perawatan terkini semakin mencabar disebabkan kemunculan Methicillin-resistant *Staphylococcus aureus* (MRSA). Kajian ini bertujuan untuk megisolasikan dan mengkategorikan MRSA yang diambil dari sistem penghawa dingin yang terdapat di Fakulti Sains dan Teknologi Sumber, UNIMAS. Sampel swab diisolasikan dari sistem ventilasi penghawa dingin sentral dan diletakkan ke dalam 0.1 % air pepton. Sampel tersebut kemudiannya dibawa untuk dijalankan ujian pencairan bersiri dan seterusnya penyaduran di atas mannitol salt agar (MSA), serta diinkubasikan selama 24-36 jam dalam suhu bilik. Hanya 16.67% daripada 24 sampel dengan faktor pencairan 10^{-5} disyaki positif *S. aureus*, dan diteruskan untuk menjalani ujian biokimia. Namun, hanya 25% daripada sampel isolasi diuji positif bagi ujian catalase, hidrolisis gelatin, methyl red, kecuali ujian sitrat dan urease. Kesemua sampel isolasi kemudian diuji ketahanannya terhadap empat antibiotik dengan menggunakan kaedah difusi disk. Kesemua sampel isolasi dianggap tahan (0 mm zon perencatan) terhadap methicillin, penisilin, vancomycin, dan cephalothin. Kajian ini penting bagi meningkatkan kesedaran staf-staf UNIMAS, pensyarah dan para pelajar dengan menyediakan data asas kepada pihak berwajib UNIMAS tentang status MRSA yang terdapat dalam sistem penghawa dingin di UNIMAS.

Kata kunci; Methicillin-resistant *Staphylococcus aureus* (MRSA), mannitol salt agar (MSA), methicillin, kaedah difusi disk.

TABLE OF CONTENTS

DECLARATION	i
ACKNOWLEDGEMENT	iii
ABSTRACT	iv
TABLE OF CONTENT	v
LIST OF TABLES	vii
LIST OF FIGURES	viii
LIST OF ABBREVIATIONS	ix
1.0 INTRODUCTION	1
2.0 LITERATURE REVIEW	4
2.1 Background of <i>Staphylococcus aureus</i>	4
2.2 Emergence of Methicillin-resistant <i>Staphylococcus aureus</i>	4
2.3 Community-acquired (CA-MRSA) and healthcare-associated (HA-MRSA).....	5
2.4 Recent studies of Methicillin-resistant <i>Staphylococcus aureus</i> in Malaysia.....	5
2.5 Disc diffusion method.....	6
2.6 Air conditioning system.....	7
2.7 Airborne bacteria.....	7
2.8 Biochemical test	
2.8.1 Catalase test.....	8
2.8.2 Methyl red test.....	8
2.8.3 Citrate test.....	8
2.8.4 Urease test.....	9
2.8.5 Gelatin hydrolysis.....	9
2.9 Antibiotic resistance	
2.9.1 Antibiotic resistance.....	10
2.9.2 Methicillin.....	10
2.9.3 Penicillin.....	11
2.9.4 Cephalothin.....	11
2.9.5 Vancomycin.....	12

3.0 MATERIALS AND METHOD	13
3.1 Swab samples collection.....	13
3.2 Determination of colony forming unit.....	15
3.3 Preliminary screening for identification of suspected <i>Staphylococcus aureus</i>	15
3.4 Disc diffusion method.....	16
4.0 RESULTS	17
5.0 DISCUSSION	25
6.0 CONCLUSION	29
7.0 RECOMMENDATIONS	30
8.0 REFERENCES	31
9.0 APPENDICES	

LIST OF TABLES

Tables	Title	Page
3.1	Detail location of the sampling	14
4.1	CFU unit per ml, colony appearance on mannitol salt agar, gram stain, and the suspected <i>S. aureus</i>	18
4.2	Results of catalase test, gelatin hydrolysis, methyl red test, citrate test, and also urease test for the samples.	22
4.3	Diameter of zone of inhibition (mm) for antibiotic resistance of methicillin, cephalothin, penicillin, and also vancomycin towards suspected MRSA.	24

LIST OF FIGURES

Figures	Title	Page
4.1	Whitish colony of suspected <i>S. aureus</i> on MSA agar	18
4.2	Morphology of suspected <i>S. aureus</i> with gram-positive (purple in colour) and have shape of cocci in clusters (100x magnification)	18
4.3	Reaction after the sample dropped with 3% H ₂ O ₂ . Immediate bubble formation (positive result on left side) and no reaction (negative result on right side)	21
4.4	Gelatin liquefaction (positive result for picture at above) and gelatin solidified (negative result for picture at bottom)	21
4.5	No changes in colour of the MRVP broth (negative result on left side) and changes in colour from yellow to red (positive result, after being dropped by methyl red reagent on right side)	22
4.6	No zone of inhibition by antibiotic (methicillin, penicillin, cephalothin, and vancomycin) on <i>S.aureus</i> that have been swabbed on Mueller-Hinton agar	24

LIST OF ABBREVIATIONS

Ammonium salts	NH ₄ H ₂ PO ₄
Beta-lactam	β-lactam
Colony forming units	CFU
Community-acquired	CA-MRSA
Healthcare-associated	HA-MRSA
Hydrogen peroxide	H ₂ O ₂
Luria Bertani broth.	LB broth
Mannitol Salt Agar	MSA
Mueller-Hinton Agar	MHA
Methicillin-resistant <i>S. aureus</i>	MRSA
Methicillin-sensitive <i>S. aureus</i>	MSSA
Microlitre	mm
Millilitre	ml
Millimetre	μl
Oxygen	O ₂
Penicillin-binding proteins	PBPs
Peptone Buffered Saline	PBS
Potential of hydrogen	pH
<i>Staphylococcus aureus</i>	<i>S. aureus</i>
Vancomycin intermediate-resistant <i>S. aureus</i>	VISA
Water	H ₂ O

1.0 INTRODUCTION

National Institute of Allergy and Infectious Diseases (2017) reported that *Staphylococcus aureus*, which normally known as staph, was found in the 1880s and in 1940s, the infection of *Staphylococcus aureus* were successfully overcome with the aids in the discovery of antibiotic, such as penicillin. But in 1961, the first strains of *S. aureus* that possessed resistance towards methicillin was discovered, which this was the first emergence of Methicillin-resistant *Staphylococcus aureus* (MRSA). Methicillin-resistant *Staphylococcus aureus* (MRSA) is nowadays the most frequently identified antibiotic-resistant pathogen in the world, including Europe, America, North Africa, the Middle East, and also East Asia. Ippolito *et al.* (2010) stated that methicillin was first introduced in 1959 in order to treat infections produced by penicillin-resistant *S. aureus*. In 1961 it was reported from United Kingdom that *S. aureus* isolates that had obtained resistance to methicillin, and since early 1970s, the occurrence of this resistance has continuously increased (Ippolito *et al.*, 2010).

MRSA is a type of *Staphylococcus aureus* which possess resistance towards antibiotics that are called beta-lactams. Beta-lactam antibiotics are among the most frequently prescribed drugs, characterized together based on their similar structural feature, which is the beta-lactam ring (Letourneau, 2017). Examples of these antibiotics include methicillin and also other typical antibiotics, for instance oxacillin, penicillin, and also amoxicillin. Majority of MRSA infections are skin infections and assumed to be non-invasive. Letourneau (2017) further said that when a person is being infected by MRSA, but it does not cause any illness towards that individual, it is called as “colonization”. According to Dani (2014), colonization defined as the presence of bacteria on a body surface without producing any disease to the individual.

Gomph (2017) explained that there are two main methods on how MRSA able to infect people. Firstly is through physical contact with infected person or a carrier (people who are not infected but are colonized with the bacteria on their body surfaces) of MRSA. The other way is for by having a physically contact MRSA from objects, for instances door handles, floors, sinks, or towels that have been touched by a MRSA-infected individual or carrier.

MRSA is hard to treat because of its resistance towards mostly of existing antibiotics. The emergence of methicillin resistance was accompanied by the development of resistance to most of the non-beta-lactam antibiotics and resulted in the reduction in options for treating infections caused by MRSA. Treatment with vancomycin, which is a glycopeptide antibiotic which always considered as a last line of defence against MRSA, has led to the emergence of vancomycin-resistant *S. aureus* (VRSA), (Rogers, 2018). In addition, Rogers (2018) added that the usage of teicoplanin, an antibiotic derived from vancomycin, also has given rise to teicoplanin-resistant MRSA strains.

MRSA causes a mild to severe infections which its infections are hardly to treat due to its resistance to several antibiotics and transmission of many virulence factors. Chew *et al.* (2018) carried out study in Malaysia to MRSA, to determine the mechanism of antibacterial activity of *Bauhinia kockiana* that arise from Peninsular Malaysia towards MRSA. Sit *et al.* (2017) reported that the prevalence rate of MRSA infection appeared to be the highest in Asia, however there is still a small number of epidemiological study carried out in Malaysia. Furthermore, there are still a poor understanding on the circumstances that mediating the ability to cause outbreaks in healthcare facilities (Boswihi & Udo, 2018).

Based on a study conducted by Shiomori *et al.* (2001), Methicillin-resistant *S. aureus* was recirculated among the patients, the air, and the inanimate environments,

especially when there was movement in the rooms. Shabha and Higgins (2011) stated that *S. aureus* is able to tolerate desiccation at a higher temperature (18°C to 37°C), and this characteristic makes it more likely to spread through ventilation and air conditioning systems.

The objectives of this study are:

- 1) To determine the presence of methicillin-resistant *Staphylococcus aureus* (MRSA) from air conditioning system in Faculty of Resource Science and Technology, UNIMAS.
- 2) To isolate the methicillin-resistant *Staphylococcus aureus* (MRSA) from air conditioning system in Faculty of Resource Science and Technology, UNIMAS successfully.
- 3) To determine the presence of methicillin-resistant *Staphylococcus aureus* (MRSA) based on biochemical and antibiotic resistance testing.

Problem statement:

The prevalence and exposure towards methicillin-resistant *Staphylococcus aureus* in air conditioning system remains unclear. As UNIMAS is now entering its 25th year of establishment, it is essential to determine if the air conditioning system do not harbour pathogenic organism. This will provide basic information to the relevant authorities and staff on the risk associated with this organism.

2.0 LITERATURE REVIEW

2.1 Background of *Staphylococcus aureus*

National Institute of Allergy and Infectious Diseases (2017) reported that *Staphylococcus aureus*, which normally known as staph, was found in the 1880s, where its infection normally created a skin condition such as boils, scalded-skin syndrome, and also impetigo, which are very hurting to the person infected. Furthermore, a more serious form of *S. aureus* infections are able to cause bacterial pneumonia and bacterial infection in our bloodstream, which both can cause mortality. In the past two decades, there were two clear shifts in the scope of epidemiology of *S. aureus* infection. Firstly, an increasing number of cases of health care associated with *S. aureus* infection, and the other one is that an epidemic of society having skin and soft tissue infections caused by *S. aureus* infection along with their resistance towards β -lactam antibiotics (Tong *et al.*, 2015).

2.2 Emergence of Methicillin-resistant *Staphylococcus aureus*

In 1940s, the infection of *Staphylococcus aureus* were successfully overcome with the aids in the discovery of antibiotic, such as penicillin. However, from that point onwards, due to its usage, as well as its overuse and misuse of antibiotics, the bacteria have been able to gain resistance to those antibiotics through evolution (National Institute of Allergy and Infectious Diseases, 2017). Therefore, methicillin, which is a form of penicillin was produced and used to counter back the infection of penicillin-resistant *S. aureus*. But in 1961, the first strains of *S. aureus* that possessed resistance towards methicillin was discovered, which this was the first emergence of Methicillin-resistant *Staphylococcus aureus* (MRSA). MRSA have been known as an essential human pathogens which caused from nosocomial infections globally (Montazeri *et al.*, 2014).

2.3 Community-acquired (CA-MRSA) and healthcare-associated (HA-MRSA)

According to Amin and Batts (2018), there are two types of *S. aureus* which are methicillin-sensitive *S aureus* (MSSA) and methicillin-resistant *S aureus* (MRSA). Under MRSA strains, there are another 2 types which are categorised as community-acquired (CA-MRSA) and healthcare-associated (HA-MRSA). Infections obtained by persons who have not been hospitalized recently (within the past year) or had a medical procedure (for instance dialysis, surgery, or catheter) are defined as CA-MRSA infections (Amin & Batts, 2018). Abigail (2008) categorised HA-MRSA as the infection in a person if the patient was in a high-risk healthcare site not long before diagnosis, or has been admitted in a facility for more than 72 hours before the infection starts to appear.

2.4 Recent studies of Methicillin-resistant *Staphylococcus aureus* in Malaysia

Soo (2016) reported that over 50% of *Staphylococcus aureus* infection are caused by Methicillin-resistant *S. aureus* (MRSA). MRSA is any type of *S. aureus* that gains resistance towards penicillins such as methicillin, oxacillin and flucloxacillin, and Malaysia alone recorded a high percentage of incidence occurred which were 32% (Soo, 2016). Over 50% of incidence were highly widespread and were reported in Asia, Malta, North and South America, and in Malaysia, the widespread of MRSA was estimated from 17% in 1986 to 44.1% in 2007 (Sit *et al.*, 2017). They further added in the journal that the prevalence rate of MRSA infection appeared to be the highest in Asia, however there is still a small number of epidemiological study carried out in Malaysia.

2.5 Disc diffusion method

Antibiotic resistance testing can be applied for discovery of drug, epidemiology, and also prognosis of medicinal result. Various of laboratory technique can be applied to assess or screen the in vitro antimicrobial action of an extract or a pure compound. One of the most popular technique is the disc-diffusion method. Balouiri *et al.* (2015) said that in this common method, agar plates are introduced with a standardized inoculum of the test microorganism. Next, 6 mm in diameter of filter paper discs that composed of desired compound at desired concentration are placed onto the surface of the agar plate.

The plate are then incubated under appropriate conditions. Normally, the antimicrobial compound diffuses into the agar and prevents development and growth of the test microorganism and then the diameters of zone of inhibition will be measured. The disk diffusion method is conducted by using Mueller-Hinton Agar (MHA), which is the most appropriate medium for regular antibiotic resistance tests due to its good reproducibility, low in sulfonamide, trimethoprim, and tetracycline inhibitors, and gives beneficial growth of most bacterial pathogens (Tendencia, 2004).

2.6 Air conditioning system

Air conditioning systems is used as a temperature cooling appliance in adjusting air temperature at home, offices, and other buildings for many purposes. It comprised with centralized component that gives an ambiance with manageable temperature, humidity, as well as purity whenever desired, and regardless of the climate state. However, these kinds of artificial environments give advantage for the growth of fungi, bacteria, protozoan and mites, which likely will cause health effect to the users, either through hypersensitivity or by infections. Anas *et al.* (2016) reported that microorganisms such as *Legionella pneumophilla*, *Streptococcus pneumonia* and *Bacillus spp* exist in air conditioner filters. In addition, study conducted by Modebelu and Modebelu (2013) had proved that air conditioner has the highest propensity to harbour and spread microorganisms if compared with other ventilation equipment that are used to provide a better air quality.

2.7 Airborne bacteria

Yassin and Almouqatea (2010) reported that when individuals get exposed to bio-aerosols that consist of airborne bacteria as well as their secondary metabolites, it can lead to respiratory problems and other health complications, for instance infections, hypersensitivity pneumonitis and also toxic response. In the air surrounding, moulds and microorganisms will become airborne and thus ubiquitous, because they able to pass through indoor areas by passive ventilation or via ventilation systems (Yassin and Almouqatea, 2010). Study conducted by Rajasekar and Balasubramaniam (2011) revealed that the dominant airborne microorganisms genera were *Staphylococcus*, *Pseudomonas*, *Alcaligenes*, and *Corynebacterium*.

2.8.1 Catalase test

Catalase is the enzyme that breaks hydrogen peroxide (H_2O_2) into H_2O and O_2 . Hydrogen peroxide is always used as a disinfectant in wounds, and the immediate bubbling formation observed is due to the production of O_2 gas. H_2O_2 is produced by some bacteria as an oxidative end product of the aerobic breakdown of sugars, and if it highly toxic to bacteria if accumulated and can lead to apoptosis (Public Health England, 2014). Since H_2O_2 is unstable, the positive and negative control should be carried out continuously.

2.8.2 Methyl red test

The Methyl-Red test tests for the potential of the bacteria to carry out mixed-acid fermentation. After incubation, the pH indicator Methyl Red is added to the broth. Methyl Red is red at pH below 4.4 (indicate positive result) and yellow at pH above 6.0. An orange colour shows an intermediate pH and would be assumed as a negative result (Hemraj, Diksha, and Avneet, 2013).

2.8.3 Citrate test

Citrate utilisation test is performed to test the capability of an organism to use sodium citrate as a sole source of carbon and ammonium salt as a sole source of nitrogen (Pradhan, 2013). Bacteria that able to grow in the medium will turn the medium alkaline. This is shown by the change of colour of bromothymol blue indicator from green to blue. Simmons Citrate agar is inoculated on the slant by touching a colony that is 18-24 hrs old with a wire loop. Inoculation from the broth culture should not be conducted since the inoculum will be too heavy. Then, it will be incubated at $35^{\circ}C$ - $37^{\circ}C$ for up to 7 days, and the blue colouration of the media is observed.

2.8.4 Urease test

According to Public Health England (2014), the urease test is performed to determine the potential of an organism to break urea, through the formation of the enzyme urease. Two units of ammonia are produced with resulting alkalinity in the presence of the enzyme, and the increased pH is detected by a pH indicator. Christensen's urea medium consists of the pH indicator phenol red which is yellow if under acid conditions (pH 6.8) and changes into rose pink in alkaline conditions (pH 8.4). The slope is inoculated heavily over the all surface and stabbed with loop wire, then it will be incubated at 35°C-37°C in an incubator. After that, the slopes will be observed after 4 hours and after overnight incubation.

2.8.5 Gelatin hydrolysis

In 1926, Frazier explained the main method to screen the ability of microorganisms to liquefy gelatin (Cruz and Torres, 2012). Thirst (1957) stated that gelatinase activity in bacteria is determined by methods which normally need the progeny of a small inoculum to liquefy relatively big amounts of substrate at a temperature unsuitable to the development of the organism. It determined the capability of bacteria to form gelatinases. It also able to differentiate the gelatinase-positive, pathogenic *Staphylococcus aureus* from the gelatinase-negative, and also non-pathogenic *Staphylococcus epidermidis*.

2.9.1 Antibiotic resistance

Physicians were finally forced to discard their belief that, given the vast array of effective antimicrobial agents, virtually all bacterial infections were treatable in the early 1970s (Lowy, 2003). Their confidence was shaken by the appearance of resistance to many antibiotics among pathogens for examples *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Pseudomonas aeruginosa*, and *Mycobacterium tuberculosis*. Foster (2017) stated that resistance can either formed by horizontal transmission of resistance determinants encoded by mobile genetic elements via plasmids, transposons, and the staphylococcal cassette chromosome, or by mutations in chromosomal genes.

2.9.2 Methicillin

In 1961, emergence of methicillin as the first of the semisynthetic penicillinase-resistant penicillin was speedily followed by reports of methicillin-resistant isolates (Lowy, 2003). Based on Stapleton and Taylor (2002), methicillin, a β -lactam antibiotic, reacts by inhibiting penicillin-binding proteins (PBPs) that are included in the formation of peptidoglycan, an important mesh-like polymer that surrounds the cell. They further explained that *Staphylococcus aureus* can become resistant to methicillin and other β -lactam antibiotics from the expression of a foreign PBP, PBP2a, that is resistant to the reaction of methicillin however still can functioning as the host PBPs (Stapleton and Taylor, 2002). Methicillin-resistant *S. aureus* isolates are always resistant to other groups of antibiotics, via variety of mechanisms which making treatment choices narrow, and this has resulted to the seek for new compounds active towards these strains.

2.9.3 Penicillin

Penicillin was introduced in 1928 but it was not utilized in clinical practice prior to 1941 (Cheng *et al.*, 2016). Widespread use of penicillin was followed by the formation of penicillin resistance among *Staphylococcus aureus* isolates within four years. Cheng *et al.* (2016) further explained that the widespread penicillin usage quickly led to the formation of penicillin resistance in *S. aureus* and new data proposed that penicillin susceptibility may be in a period of renaissance. (Cheng *et al.*, 2016). According to Hagstrand, Skovby, and Pahlman (2017), penicillin is now seldom used in the clinical practices of *S. aureus* infections due to widespread resistance and a worry about the exactness of existing techniques for penicillin susceptibility testing.

2.9.4 Cephalothin

Cephalothin is a first generation cephalosporin antibiotic and remains to be widely used (Choudhary, 2017). Cephalosporins are beta-lactam antimicrobials that have same mechanisms of action and a similar structure with penicillins. Harrison and Bratcher (2008) explained that penicillins and cephalosporins possess the same four-member “core” beta-lactam ring, however cephalosporins have an extra atom in the side ring. Mechanism of action of cephalosporins consist of a rigid bacterial cell walls that can be assumed as a series of repeating interlinking units reminiscent of floor tiles (Harrison and Bratcher, 2008). During replication, a bacterium discards “tiles” circumferentially to let cell division through a pinching-like action, while rapidly substituting new “tiles” at the ends of what have become two bacteria. This process needs enzymes to interlink replacement tiles. Such enzymes are the aims of beta-lactam antibiotics and are known as penicillin-binding proteins (PBPs). Antibiotic process requires binding to PBPs, averting them from closing the vulnerable ends on dividing bacteria and resulting the natural intrabacterial hyperosmotic pressure to break down the bacteria. Therefore, beta-lactam antibiotics are bactericidal.

2.9.5 Vancomycin

Methicillin-resistant *Staphylococcus aureus* probably sensitive to some other antibiotics, such as clindamycin, macrolides, tetracyclines, trimethoprim-sulfamethoxazole and quinolones, or resistant to all antibiotics except vancomycin (Hakim *et al.*, 2007). Hakim *et al.* (2007) explained that vancomycin stayed as the only predictable active antibiotic towards all strains of *S. aureus*, and especially methicillin-resistance *S. aureus*. Vancomycin, a glycopeptide antibiotic that obstructs cell wall biosynthesis, stays as a best choice of drug for treatment of serious MRSA infections. According to McGuinness *et al.* (2017), *S. aureus* strains showing increased resistance towards vancomycin, which called vancomycin intermediate-resistant *S. aureus* (VISA) were first known in the 1990s. The molecular basis of resistance in VISA is polygenic and involves stepwise mutations in genes encoding molecules predominantly included in cell envelope biosynthesis (McGuinness *et al.*, 2017). *S. aureus* isolates with fully resistance towards vancomycin are called as vancomycin-resistant *S. aureus* (VRSA), and they were first reported in the U.S. in year 2002.

3.0 MATERIALS AND METHODS

3.1 Swab samples collection

A total of 24 swab samples were collected from 12 lecturers' offices, with each of the offices were chosen as the representative of the main office based on its location. The offices are located in Faculty of Resource Science and Technology, UNIMAS. The swab samples were collected from the evaporator of the central air conditioning system. The cotton bud was dipped into a sterile water prior to swabbing. After the swab samples collected, it was inserted into the Bijou bottle, till the bottom of the medium of 3 ml of 0.1 % peptone water, and were kept in icebox. The swab samples were then transported to the Microbiology laboratory as soon as possible, and were proceeded to serial dilution. Table 3.1 shows the detail location of the samplings taken. Two offices were chosen as the representative at each location, and all of the swab samples were collected from central air conditioning system.