



**Citation:** Wong Sin Yeng, Peter C. Boyce (2023) Homalomeneae (Araceae) of Borneo XXX — *Homalomena atlas* [Hanneae Complex] a new species from the Rejang Basin, Sarawak. *Webbia. Journal of Plant Taxonomy and Geography* 78(1): 33-37. doi: 10.36253/jopt-14254

**Received:** Jan 31, 2023

**Accepted:** Feb 9, 2023

**Published:** May 14, 2023

**Copyright:** ©2023 Wong Sin Yeng, Peter C. Boyce. This is an open access, peer-reviewed article published by Firenze University Press (<http://www.fupress.com/webbia>) and distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Competing Interests:** The Author(s) declare(s) no conflict of interest.

**Editor:** Alistair Hay

**ORCID**

WSY: 0000-0003-4042-9672

PCB: 0000-0002-5856-9561

## Homalomeneae (Araceae) of Borneo XXX — *Homalomena atlas* [Hanneae Complex] a new species from the Rejang Basin, Sarawak

WONG SIN YENG<sup>1,2,\*</sup>, PETER C. BOYCE<sup>3</sup>

<sup>1</sup> Institute of Biodiversity and Environmental Conservation, Universiti Malaysia Sarawak 94300 Kota Samarahan, Sarawak, Malaysia

<sup>2</sup> Harvard University Herbaria, Cambridge, MA, USA.

<sup>3</sup> Dipartimento di Biologia, Centro Studi Erbario Tropicale (herbarium FT), University of Florence, Via G. La Pira 4, 50121, Firenze, Italy

\*Corresponding author. E-mail: sywong@unimas.my

**Abstract.** *Homalomena atlas* is described and illustrated as a new species from the western Rejang Basin and assigned to the Hanneae Complex of the Cyrtocladon Clade, taking the complex to 12 described species on Borneo.

**Keywords:** Araceae, Borneo, Cyrtocladon Clade, Hanneae Complex, *Homalomena* Sarawak, Shales.

### INTRODUCTION

The *Homalomena* Hanneae Complex (Ng et al. 2011: 26) of the Cyrtocladon Clade (Wong et al. 2013) is a highly distinctive group of species notable for leaf blades with conspicuous extrafloral nectaries, amber-coloured resin secreted from between the staminate florets shortly before staminate anthesis (Hoe et al. 2011, 2016; Wong & Boyce 2017), and spathes white at anthesis with the persistent post-anthesis spathe turning deep pink to red as the fruits develop. Until recognition of *Homalomena electra* P.C.Boyce & S.Y.Wong (Boyce & Wong 2017) from Peninsular Malaysia the Hanneae Complex was considered to be entirely Bornean (but see Wong et al. 2013: 592). Including that described here, the complex comprises 13 described species, all except *H. electra* restricted to Borneo, and with most species localized. Based on field observations and in particular the huge amount of undeterminable material deposited in herbaria the Hanneae Complex is most probably contains well in excess of 100 species

Geology in this paper is specified based on Hutchinson (1989, 2005) and Tate (2001).