

# Modern human DNA analyses with special reference to the inner dual-structure model of Yaponesian

Timothy A. JINAM<sup>1,2</sup>, Yosuke KAWAI<sup>3</sup>, Naruya SAITOU<sup>1,2,4,5\*</sup>

<sup>1</sup>Population Genetics Laboratory, National Institute of Genetics, Mishima City, Shizuoka 411-8540, Japan

<sup>2</sup>Department of Genetics, School of Life Science, Graduate University for Advanced Studies (SOKENDAI), Mishima City, Shizuoka 411-8540, Japan

<sup>3</sup>Genome Medical Science Project, National Center for Global Health and Medicine, Toyama, Shinjuku-ku, Tokyo 162-8655, Japan

<sup>4</sup>Faculty of Medicine, University of The Ryukyus, Nishihara-cho, Okinawa 903-0215, Japan

<sup>5</sup>Department of Biological Sciences, Graduate School of Science, the University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan

Received 2 October 2020; accepted 17 December 2020

**Abstract** Previous studies suggested two major migration events during the Jomon and Yayoi periods that affected the genetic diversity of modern Japanese (Yaponesian). We explored the possibility of a three-wave migration model by examining three datasets of modern human DNA: (1) whole mitochondrial (mt) DNA genomes of 1642 Yaponesians; (2) mtDNA haplogroup frequencies of 59105 Yaponesians from 47 prefectures; and (3) genome-wide SNP data of two Yaponesians (Ainu, Okinawa) and whole-genome sequence data of Yamato individuals, the Funadomari Jomon F23 individual, and three East Asian populations (Korean, northern Chinese, and southern Chinese). Past population size change was estimated based on dataset 1, and we clearly observed a steep population increase after the Yayoi period. Principal-component analysis and phylogenetic network analysis were applied to dataset 2, and we confirmed the pattern consistent with our model. An admixture program was used on dataset 3, and we found that the two- and three-layer migration models are both compatible with these SNP data. Taken together, these three datasets provide support for our three-wave, ‘inner dual-structure’ model.

**Key words:** Yaponesia Genome Project, mitochondrial DNA, population size changes, three-wave migration model, inner dual-structure model

## Introduction

The term ‘Yaponesia’ was proposed by Toshio Shimao in the 1960s (e.g. Shimao, 1977) to refer to the Japanese archipelago, and later Saitou (2015, 2017) defined the three geographical areas of Yaponesia: northern Yaponesia (Sakhalin Island, Kuril Islands, and Hokkaido Island); central Yaponesia (Honshu, Shikoku, Kyushu Islands, and surrounding smaller islands); and southern Yaponesia (Nansei Islands including Amami and Okinawa regions). Saitou (2015, 2017) and Saitou and Jinam (2017) proposed a three-wave migration model for Yaponesia. The estimated time frames of these migrations according to Saitou (2017) are: (1) Paleolithic and the middle period of Jomon (40000 years before present (BP) to 4500 BP) for the first wave; (2) late and final Jomon periods (4500 BP to 3000 BP) for the second wave; (3) Yayoi period (3000 BP) to present day for the third wave.

The people involved in those three waves may be hypoth-

esized as follows. Hunter-gatherers from Siberia, continental East Asia, and/or Indochina migrated to Yaponesia during the first wave. This migration wave has been well discussed (e.g. Suzuki, 1983; Hanihara, 1991; Omoto and Saitou, 1997). We propose a somewhat mysterious ‘sea people’ as the second-wave migrants. They may have been hunter-gatherers who mainly relied on fishing and were distributed from the coastal area of southern China to the Shandong peninsula, the Yellow Sea, and the Korean peninsula. Shinoda et al. (2019) analyzed ancient DNA of people who lived on Gadeok Island, located of the southeast coast of Korean peninsula about 6300 years ago, and reported that they had a much higher Jomon component than modern Koreans. Shima (2020) recently stressed the importance of fishing in modern human dispersal. These two literatures seem to support the existence of ancient hunter-gatherers who mainly relied on fishing on coastal areas of continental East Asia. These ‘sea people’ were replaced by rice farmers whose population size quickly expanded between 7000 BP and 5000 BP, and these ‘sea people’ eventually migrated to Yaponesia during the late and final Jomon periods (4500 BP–3000 BP). There is a possibility that these ‘sea people’ spoke an ancient Japanese language, which could have been the ‘lingua franca’ of people in coastal East Asia at that time (Saitou, 2015, 2017). Rice agriculture was introduced c.

\* Correspondence to: Naruya Saitou, Population Genetics Laboratory, National Institute of Genetics, Mishima City, Shizuoka 411-8540, Japan.

E-mail: [saitounr@nig.ac.jp](mailto:saitounr@nig.ac.jp)

Published online 31 March 2021

in J-STAGE ([www.jstage.jst.go.jp](http://www.jstage.jst.go.jp)) DOI: 10.1537/ase.201217

Table 1. List of complete mtDNA genomes used

Sample description	Reference	No. of individuals
Published data of Japanese:		
Healthy individuals from Tokyo and Nagoya	Tanaka et al. (2004)	657
Atypical psychosis patients	Kazuno et al. (2005)	57
Semi-supercentenarians	Bilal et al. (2008)	112
Schizophrenic patients	Ueno et al. (2009)	90
Healthy unrelated people	Nohira et al. (2010)	14
Athletes	Mikami et al. (2013)	185
Human Genome Diversity Project (HGDP)	Bergström et al. (2020) and 1 <sup>a</sup>	27
Simons Genome Diversity Project (SGDP)	Mallick et al. (2016) and 2 <sup>a</sup>	3
Unpublished data of Japanese:		
People living in Tokyo area	Nishida et al. (2008)	417
People originating from Izumo area	Saitou et al. (unpublished)	21
People originating from Makurazaki area	Saitou et al. (unpublished)	59
Subtotal (Japanese)		1642

<sup>a</sup> 1, [www.internationalgenome.org/data-portal/population/JapaneseHGDP](http://www.internationalgenome.org/data-portal/population/JapaneseHGDP); 2, [www.internationalgenome.org/data-portal/population/JapaneseSGDP](http://www.internationalgenome.org/data-portal/population/JapaneseSGDP)

3000 BP (e.g. Fujio, 2015; Nasu and Momohara, 2016) by the third-wave migrants whose homeland was somewhere in continental East Asia. This migration wave still continues today, for the highest proportion of international marriages in Japan includes Chinese and Koreans (e-Stat, 2018). We will discuss the plausibility of this three-wave migration model based on genome-wide single-nucleotide polymorphism (SNP) and mitochondrial (mt) DNA analyses.

## Materials and Methods

We utilized three kinds of datasets, each with their corresponding methods. The first dataset was a total of 1642 complete mitochondrial genomes from Yaponeseans (Table 1). These included 1115 published sequences (Tanaka et al., 2004; Kazuno et al., 2005; Bilal et al., 2008; Ueno et al., 2009; Nohira et al., 2010; Mikami et al., 2013), 417 unpublished mitochondrial genomes of people from Tokyo (DNA data are the same as is Nishida et al., 2008), and a total of 80 unpublished sequences we determined for Yaponeseans from Izumo and Makurazaki. It should be noted that 13 mtDNA sequences reported by Tanaka et al. (2004) were identified as artificial recombinants (Kong et al., 2008), so we used mtDNA data of 657 individuals, not 670 as originally reported by Tanaka et al. (2004). These 1642 Yaponesean mtDNA genome sequences were used to estimate effective population size changes over time. These sequence data were aligned using MAFFT (Katoh and Standley, 2013), and insertions and deletions were omitted. BEAST software (Drummond and Rambaut, 2007) was used to generate a Bayesian Skyline plot using two datasets: whole mtDNA (16570 bp) and a coding region (a concatenation of 13 protein-coding gene sequences; 11341 bp). The substitution model of Tamura and Nei (1993) was used, assuming a strict molecular clock with separate substitution rates for whole mtDNA and coding regions:  $2.67 \times 10^{-8}$  substitutions/site/year and  $1.57 \times 10^{-8}$  substitutions/site/year, respectively (Fu et al., 2013).

The second dataset used was mtDNA haplogroup fre-

quency data of 59105 Yaponeseans living in all 47 prefectures provided by Genesis-HealthCare, Ltd, as shown in Supplementary Table. Principal-component analysis (PCA) was performed using R version 3.6.3, Nei's (1972) standard genetic distance was calculated using phylip version 3.698 (Felsenstein, 2005), and the genetic distance matrix was used to making neighbor-net networks (Bryant and Moulton, 2004; Huson and Bryant, 2006). We classified 47 prefectures into a central axis area and a peripheral area based on Saitou's (2017) proposal as shown in Table 2 and Figure 1. The central axis of Yaponesia starts from northern Kyushu and stretches to Edo/Tokyo, the city which has been the political center of Yaponesia from the early 17th century. Northern Kyushu was the first region to adopt wet rice agriculture (e.g. Fujio, 2015) and was the political center of western Yaponesia during the early and middle Yayoi periods. The Nara-Kyoto area was the political center of Yaponesia from the late Yayoi period to the end of 16th century (also known as the Azuchi-Momoyama period in historical literature).

The third dataset comprised 639912 autosomal SNP data from two Japanese populations (Ainu, Ryukyuan) reported by the Japanese Archipelago Human Population Genetics Consortium (2012). This SNP dataset was merged with whole-genome sequence data of the Funadomari Jomon F23 individual (Kanzawa-Kiryama et al., 2019), 27 Japanese (Bergström et al., 2020), 40 Koreans (Zhang et al., 2014), 45 Han Chinese Beijing (CHB), and 45 southern Han Chinese (CHS) (Bergström et al., 2020). After pruning the dataset for linkage disequilibrium (LD), the resulting 192898 SNP data were used for admixture analysis (Alexander et al., 2009). Merging of datasets and LD pruning was performed using PLINK version 1.9 (Chang et al., 2015).

## Results and Discussion

### Dataset 1

We first estimated population size changes over time using 1642 complete mitochondrial genomes of current Japanese. There were sharp increases in the population size dur-

Table 2. Classification of 47 prefectures of Japan into two areas

Prefecture name	Abbreviation	Region
(A) 20 prefectures in central axis area (red in Figure 2)		
Aichi	AIC	Tokai
Chiba	CHI	Kanto
Fukuoka	FKK	Kyushu
Gifu	GIF	Tokai
Gunma	GUN	Kanto
Hiroshima	HIR	San-yo
Hyogo	HYO	Kinki
Ibaraki	IBA	Kanto
Kanagawa	KAN	Kanto
Kyoto	KYO	Kinki
Mie	MIE	Tokai
Nara	NAR	Kinki
Okayama	OKA	San-yo
Osaka	OSA	Kinki
Saitama	SAI	Kanto
Shiga	SGA	Kinki
Shizuoka	SZK	Tokai
Tochigi	TTG	Kanto
Tokyo	TKY	Kanto
Yamaguchi	YMG	San-yo
(B) 27 prefectures in peripheral area (white in Figure 2 except*)		
Akita	AKI	Tohoku
Aomori	AOM	Tohoku
Ehime	EHI	Shikoku
Fukui	FKI	Hokuriku
Fukushima	FKS	Tohoku
Hokkaido	HOK	Hokkaido*
Ishikawa	ISI	Hokuriku
Iwate	IWA	Tohoku
Kagawa	KGW	Shikoku
Kagoshima	KGS	Kyushu
Kochi	KOC	Shikoku
Kumamoto	KUM	Kyushu
Miyagi	MIG	Tohoku
Miyazaki	MIZ	Kyushu
Nagano	NGN	Koshin-etsu
Nagasaki	NGS	Kyushu
Niigata	NII	Koshin-etsu
Oita	OIT	Kyushu
Okinawa	OKI	Okinawa*
Saga	SAG	Kyushu
Shimane	SMN	San-in
Tokushima	TKS	Shikoku
Tottori	TTT	San-in
Toyama	TYM	Hokuriku
Wakayama	WAK	Kinki
Yamagata	YGA	Tohoku
Yamanashi	YMN	Koshin-etsu




Figure 1. Classification of 47 prefectures in Japan as central axis (red) or peripheral (white). The map was taken from WANtaroHP ([http://civilyarou.web.fc2.com/WANtaroHP\\_F90\\_html5/index.html](http://civilyarou.web.fc2.com/WANtaroHP_F90_html5/index.html)).

riod peaked during the middle Jomon (5500 BP–4500 BP), based on archaeological data (Koyama, 1978). However, this peak in population size was not observed in Figure 2A and B. Instead, much older dates (25000 BP and 38000 BP for Figure 2A and B, respectively) showed second points of population size increase, corresponding to the Paleolithic period. Saitou (2015) showed a similar result based on 1057 complete mtDNA of Japanese, and a steep population size increase after 3000 BP was also observed. However, its population dynamics was somewhat different.

Okada et al. (2018) estimated past population size changes of modern Japanese using whole nuclear DNA genome data. Their results based on 1276 complete genomes (Figure 2 in their paper) showed that the start of population size increase in the Japanese was 6000 BP, much earlier than the start of the Yayoi period, even before the middle Jomon. This time estimate was based on a “per-generation mutation rate at  $1.25 \times 10^{-8}$  mutations per base pair and a constant generation time of 29 years” (Okada et al., 2018). Harris (2015) suggested a recent change in mutation rate in human evolution. We have to be careful about time estimates, although the relative pattern of the population size changes may be robust. The bottleneck followed by a rapid increase of the population size was noted both by our Figure 2 and by Figure 2 of Okada et al. (2018). This consistency may mean that the population size increased after introduction of wet rice agriculture in the Yayoi period.

Watanabe et al. (2019) estimated the population size changes of modern Japanese males using 122 Y chromosome D haplogroup (clade 1 in their term) sequence data.

ing the last 3000 years when using either coding region only (Figure 2A) or complete mtDNA (Figure 2B). This increase in population size corresponds to the start of the Yayoi period, just after the Jomon period (16500 BP–3000 BP) (Fujio, 2015; Yamada, 2019). Population size during the Jomon pe-




Figure 2. Estimate of the Japanese population size change based on 1542 mtDNA sequence data.

They observed a clear bottleneck during the early Yayoi period followed by a sharp increase in population size. The D haplogroup is known to be distributed only among Japanese, Tibetan, and Andamanese, and is thought to have contributed to the formation of the Neolithic Jomon males in prehistoric Japan (Kanzawa-Kiriyama et al., 2019). Watanabe et al. (2019) argued that this sharp decrease was due to a shortage of food resources in the colder final Jomon period.

**Dataset 2**

Figure 6 shows a two-dimensional PCA plot of 47 Japanese prefectures using the frequencies of 36 mtDNA haplotypes frequencies presented in Supplementary Table. PC1 and PC2 explain 89% and 6.7% of the total variance, respectively. Prefectures are color-coded into two areas according to Supplementary Table and Figure 1—red for the central axis of Yaponesia and black for the peripheral area. Because Okinawa Prefecture (OKI) was so distant from the remaining 46 prefectures, the middle part of the plot was truncated for clarity. Interestingly, central axis prefectures tend to locate on the right side of Figure 3, while peripheral area prefectures are on the left side. If the 47 prefectures are equally separated along PC1 (23 and 24 on the right and left, respectively; indicated by a thin blue line in Figure 3), this biased distribution is weakly statistically significant (5% level using Fisher’s exact test applied to the 2 × 2 contingency table) compared to a random distribution, as shown in Table 3A.

Because of the unique location of Okinawa Prefecture in Figure 3, we eliminated Okinawa and conducted PCA for the remaining 46 prefectures, shown in Figure 4, with PC1 and PC2 explaining 56% and 22% of the total variance, respectively. Now Yamaguchi Prefecture (YMG, classified as




Figure 3. PCA plots of 47 prefectures based on mtDNA haplotype frequencies.

Table 3. Contingency test (2 × 2) for PCA plots of Figure 3 and Figure 4

(A) Comparison of 47 prefectures (see Figure 3)

	Left side	Right side	Sum
Central axis	6	14	20
Peripheral	18	9	27
Sum	24	23	47

*P* = 0.0189 (both sides: Fisher’s exact test)

(B) Comparison of 46 prefectures<sup>a</sup> (see Figure 4)

	Left side	Right side	Sum
Central axis	6	14	20
Peripheral	17	9	26
Sum	23	23	46

*P* = 0.0361 (both sides: Fisher’s exact test)

<sup>a</sup> Okinawa Prefecture not included.

central axis) is located at the rightmost position, within proximity to some prefectures in the Kyushu region, such as Nagasaki (NGS) and Kumamoto (KUM). Generally speaking, as in Figure 3, central-axis prefectures tend to locate on the right side and peripheral-area prefectures on the left side. Similar to Figure 3, when equally divided along PC1, the biased distribution of prefectures is again weakly statistically significant (5% level using Fisher’s exact test applied to the 2 × 2 contingency table) compared to a random distribution as shown in Table 3B.

Previously, Saitou (2017) presented a PCA plot using mtDNA haplogroup data of 47 prefectures from a total of 18641 Yaponesian individuals, also provided by Genesis-HealthCare, Ltd. Although the relative positions of prefectures slightly differed between Saitou (2017) and our current

results (Figure 3, Figure 4), the central axis–peripheral difference of Saitou (2017) was also statistically significant at the 2% level using Fisher’s exact test applied to the 2 × 2 contingency table (see also Jinam et al., 2021). The analysis shown in this paper is consistent with this pattern.

We constructed phylogenetic networks of prefectures using mtDNA haplogroup frequency data. Figure 5 shows a network of all 47 prefectures. As in Figure 3, Okinawa Prefecture is very distantly related to the remaining 46 prefectures. We thus made another phylogenetic network for the 46 prefectures without Okinawa (Figure 6). Central-axis and peripheral prefectures are colored in red and blue, respectively, in both figures. Although there are some topological differences between the two networks shown in Figure 5 and Figure 6, the overall pattern is similar. We thus explain the network of Figure 6, to have an easier grasp of the relationship between prefectures. Generally speaking, central-axis prefectures tend to locate at the center and peripheral ones are literally located peripherally. Prefectures in Kyushu (SAG, KUM, NGS, KGS, MIZ, FKK, and OIT), which are the closest to Okinawa Prefecture in Figure 5, are located in the left side of this network, while those in Tohoku (AOM, IWA, AKI, FKS, YMM, and MIG) are located in the right side. In contrast, prefectures in Shikoku (TKS, KOC, EHI, and KGW) are scattered. KOC (Kochi Prefecture) and EHI (Ehime Prefecture) formed one cluster, whereas KGW (Kagawa Prefecture) is closest to FKI (Fukui Prefecture) in Hokuriku. TKS (Tokushima Prefecture) shares a split with YMG (Yamaguchi Prefecture) in San-yo. Another interesting feature of this phylogenetic network is that prefectures in San-in (SMN and TTT), which formed one cluster, are closest to OIT (Oita Prefecture) in Kyushu. MIE (Mie Prefecture) in Kinki, SZK (Shizuoka Prefecture) in Tokai, and IBA (Ibaraki Prefecture) in Kanto are tightly clustered with a




Figure 4. PCA plots of 46 prefectures (Okinawa not included) based on mtDNA haplotype frequencies.






Figure 5. Phylogenetic network of 47 prefectures based on mtDNA haplotype frequencies.




Figure 6. Phylogenetic network of 46 prefectures (Okinawa not included) based on mtDNA haplotype frequencies.




Figure 7. (a) Admixture result from  $k = 2$  to  $k = 4$ . b) Cross-validation error of admixture from  $k = 2$  to  $k = 7$ .

long split; however, this clustering is inconsistent with their geographical locations.

Recently Watanabe et al. (2020) reported analysis of ~140000 nuclear SNP data for 11069 individuals distributed in all the 47 prefectures. They also found a clear difference between Okinawa and the remaining 46 prefectures, and all seven prefectures in Kyushu are closest to Okinawa. This pattern is consistent with that of our Figure 5.

### Dataset 3

We now used genome-wide SNP data (see Materials and Methods) for two Yaponesian populations (Ainu and Okinawa), and genome data for Jomon F23 and Yamato, and three continental East Asians (Koreans, CHB, and CHS) to estimate their ancestral population structure. Figure 7a shows the result of admixture analysis when the number of ancestral components ( $k$ ) was assumed to be 2–7. When  $k = 2$ , the two ancestral components are color-coded blue and orange. Interestingly, some Ainu individuals are 100% blue component as in the F23 Jomon individual. The lowest blue component for Ainu individuals was more than 50%. Whereas all Okinawa individuals have more or less 30% blue component, Yamato individuals are on average ~15% blue component. Among continental populations, only Korean has ~5% blue component, and CHB and CHS contain very small proportions of the blue component, if any. This admixture result for  $k = 2$  is very similar to Supplementary Figure 2 of Jinam et al. (2015), and fits well with the dual-structure model of Yaponesian formation; the blue component is indigenous (Jomon) component, and the orange component is migrants to Yaponesia after the Yayoi period.

When  $k = 3$ , the orange-colored ancestral component for  $k = 2$  was, roughly speaking, separated into green and or-

ange components, although proportions of the blue component were also slightly modified. Now most of Okinawa individuals are composed of ~90% green and ~10% blue components, and the green component is also dominant (~65%) in Yamato individuals, with orange (~30%) and blue (~5%) components. In contrast, the orange component is dominant (~70%) in Korean and the rest (~30%) were the green component, with no blue component. CHB contains 0–5% green component depending on individuals, and the orange component is 95–100%. CHS is more extreme—almost all individuals have ~100% orange component. If we fit this  $k = 3$  situation to the three-wave migration model proposed by Saitou (2015, 2017), the blue, green, and orange components may correspond to first, second, and third waves, respectively.

When  $k = 4$ , the red component appears only in continental populations, and this component is especially dominant (70–100%) among CHS individuals. The red component is also dominant in CHB; 50–100% depending on individuals. Among Koreans, however, only a small proportion of the red component exists in some individuals. If we compare cross-validation (CV) errors for different  $k$  values (Figure 7b), the fits for  $k = 4$ –7 are as bad as for  $k = 1$  (only one ancestral component), and it seems only cases when  $k = 2$  or  $k = 3$  may be considered to be more realistic.

### Conclusion

We used three datasets in this study, and each dataset was analyzed using different methods. Dataset 1 comprises mtDNA genome sequences, and the population size change in the past was inferred from these data of modern Yaponesian. These mtDNA data clearly showed a steep increase in

the population size after the Yayoi period. This confirms the shift from hunting and gathering to agriculture after the Yayoi period started. Dataset 2 comprises mtDNA haplogroup frequency data of 47 prefectures, and the dichotomy between central-axis and peripheral populations supports our ‘inner dual-structure’ model. Dataset 3 (genome-wide SNP data) seems to be consistent with the three-wave migration model. We need to compare nuclear DNA data of many areas of Yaponesia to strengthen the existence of the ‘inner dual-structure’ and the three-wave migration model.

### Acknowledgments

We thank the Asian DNA Repository Consortium for the use of genome-wide SNP data of three Japanese populations. We also thank Genesis Healthcare for providing us with mtDNA haplogroup frequency data of Japanese. Preparation of figures by Mrs. Masako Mizuguchi and Mrs. Takayo Hamasuna is appreciated. Finally, we appreciate people in Yaponesia and other geographical areas who kindly gave their DNA to the scientific community.

This paper is a part of the activity of MEXT Grant-in-Aid for Scientific Research on Innovative Areas ‘Yaponesia Genome’ ([www.yaponesian.jp](http://www.yaponesian.jp); see also Yaponesian Genome Project Managing Group, 2020).

### References

- Alexander D.H., Novembre J., and Lange K. (2009) Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, 19: 1655–1664.
- Bergström A., McCarthy S.A., Hui R., Almarri M.A., Ayub Q., et al. (2020) Insights into human genetic variation and population history from 929 diverse genomes. *Science*, 367(6484): eaay5012.
- Bilal E., Rabadan R., Alexe G., Fuku N., Ueno H., et al. (2008) Mitochondrial DNA haplogroup D4a is a marker for extreme longevity in Japan. *PLoS One*, 3: e2421.
- Bryant D. and Moulton V. (2004) Neighbor-net: an agglomerative method for the construction of phylogenetic networks. *Molecular Biology and Evolution*, 21: 255–265.
- Chang C.C., Chow C.C., Tellier L.C.A.M., Vattikuti S., Purcell S.M., and Lee J.J. (2015) Second-generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience*, 4: doi: 10.1186/s13742-015-0047-8.
- Drummond A.J. and Rambaut A. (2007) BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology*, 7: 214.
- e-Stat (2018) Table 9–18: Numbers and percentages of annual marriages based on nationalities of wives and husbands [in Japanese]: <https://www.e-stat.go.jp/dbview?sid=0003214861>
- Felsenstein J. (2005) PHYLIP (Phylogeny Inference Package) version 3.6. Distributed by the author. Department of Genome Sciences, University of Washington, Seattle, WA.
- Fu Q., Mittnik A., Johnson P.L.F., Bos K., Lari M., et al. (2013). A revised timescale for human evolution based on ancient mitochondrial genomes. *Current Biology*, 23: 553–559.
- Fujio S. (2015) History of the Yayoi Period. Kodansha, Tokyo (in Japanese).
- Hanihara K. (1991) Dual structure model for the population history of Japanese. *Japan Review*, 2: 1–33.
- Harris K. (2015) Evidence for recent, population-specific evolution of the human mutation rate. *Proceedings of the National Academy of Sciences of the United States of America*, 112: 3439–3444.
- Huson D.H. and Bryant D. (2006) Application of phylogenetic networks in evolutionary studies. *Molecular Biology and Evolution*, 23: 254–267.
- Japanese Archipelago Human Population Genetics Consortium: Jinam T., Nishida N., Hirai M., Kawamura S., Oota H., et al. (2012) The history of human populations in the Japanese archipelago inferred from genome-wide SNP data with a special reference to the Ainu and the Ryukyuan populations. *Journal of Human Genetics*, 57: 787–795.
- Jinam T.A., Kanzawa-Kiriyama H., Inoue I., Tokunaga K., Omoto K., and Saitou N. (2015) Unique characteristics of the Ainu population in northern Japan. *Journal of Human Genetics*, 60: 565–571.
- Jinam T.A., Kawai Y., Kamatani Y., Sonoda S., Makisumi K., et al. (2021) Genome-wide SNP data of Izumo and Makurazaki populations support inner-dual structure model for origin of Yamato people. *Journal of Human Genetics* (in press).
- Kanzawa-Kiriyama H., Jinam T.A., Kawai Y., Sato T., Hosomichi K., et al. (2019) Late Jomon male and female genome sequences from the Funadomari site in Hokkaido, Japan. *Anthropological Science*, 127: 83–108.
- Katoh K. and Standley D.M. (2013) MAFFT Multiple Sequence Alignment Software version 7: improvements in performance and usability. *Molecular Biology and Evolution*, 30: 772–780.
- Kazuno A., Munakata K., Mori K., Tanaka M., Nanko S., et al. (2005) Mitochondrial DNA sequence analysis of patients with ‘atypical psychosis.’ *Psychiatry and Clinical Neurosciences*, 59(4): 497–503.
- Kong Q.P., Salas A., Sun C., Fuku N., Tanaka M., et al. (2008) Distilling artificial recombinants from large sets of complete mtDNA genomes. *PLoS One*, 3: e3016.
- Koyama S. (1978) Jomon subsistence and population. *Senri Ethnological Studies*, 2: 1–65.
- Mallick S., Li H., Lipson M., Mathieson I., Gymrek M., Racimo F., et al. (2016) The Simons Genome Diversity Project: 300 genomes from 142 diverse populations. *Nature* 538(7624): 201–206.
- Mikami E., Fuku N., Kong Q.-P., Takahashi H., Ohiwa N., et al. (2013) Comprehensive analysis of common and rare mitochondrial DNA variants in elite Japanese athletes: a case-control study. *Journal of Human Genetics*, 58: 780–787.
- Nasu H. and Momohara A. (2016) The beginnings of rice and millet agriculture in prehistoric Japan. *Quaternary International*, 397: 504–512.
- Nei M. (1972) Genetic distance between populations. *American Naturalist*, 106: 283–292.
- Nishida N., Koike A., Tajima A., Ogasawara Y., Ishibashi Y., Uehara Y., et al. (2008) Evaluating the performance of Affymetrix SNP Array 6.0 platform with 400 Japanese individuals. *BMC Genomics*, 9: 431.
- Nohira C., Maruyama S., and Minaguchi, K. (2010) Phylogenetic classification of Japanese mtDNA assisted by complete mitochondrial DNA sequences. *International Journal of Legal Medicine*, 124(1): 7.
- Okada Y., Momozawa Y., Sakaue S., Kanai M., Ishigaki K., et al. (2018) Deep whole-genome sequencing reveals recent selection signatures linked to evolution and disease risk of Japanese. *Nature Communications*, 9: 1631.
- Omoto K. and Saitou N. (1997) Genetic origins of the Japanese: a partial support for the dual structure hypothesis. *American Journal of Physical Anthropology*, 102: 437–446.
- Saitou N. (2015) History of People on the Japanese Archipelago. Iwanami Shoten, Tokyo (in Japanese).
- Saitou N. (2017) Origin of Japanese Viewed from Nuclear DNA Analyses. Kawade Shobo Shinsha, Tokyo (in Japanese).
- Saitou N. and Jinam T.A. (2017) Language diversity of the Japanese archipelago and its relationship with human DNA diversity. *Man in India*, 97: 205–228.
- Shima T. (2020) Human History of Fish Eating—From Out-of-Africa to Japanese Archipelago. NHK Shuppan, Tokyo (in



- Japanese).
- Shimao T. (1977). *Thinking Yaponesia*. Ashi Shobo, Tokyo (in Japanese).
- Shinoda K., Kanzawa-Kiriyama H., Kakuda T., and Adachi N. (2019) Ancient DNA analysis of skeletal remains excavated from the Jang Hang site in Gadeokdo Islands, Busan-city, Korea (in Japanese and in Korean). *Journal of MUNMUL*, 9: 167–206.
- Suzuki H. (1983) *Roots of Japanese Viewed from Bones*. Iwanami Shinsho. Iwanami Shoten, Tokyo (in Japanese).
- Tamura K. and Nei M. (1993) Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution*, 10: 512–526.
- Tanaka M., Cabrera V.M., González A.M., Larruga J.M., Takeyasu T., et al. (2004) Mitochondrial genome variation in Eastern Asia and the peopling of Japan. *Genome Research*, 14: 1832–1850.
- Ueno H., Nishigaki Y., Kong Q.-P., Fuku N., Kojima S., et al. (2009). Analysis of mitochondrial DNA variants in Japanese patients with schizophrenia. *Mitochondrion*, 9: 385–393.
- Watanabe Y., Naka I., Khor S.-S., Sawai H., Hitomi Y., et al. (2019) Analysis of whole Y-chromosome sequences reveals the Japanese population history in the Jomon period. *Scientific Reports*, 9: 8556.
- Watanabe Y., Isshiki M., and Ohashi J. (2020) Prefecture-level population structure of the Japanese based on SNP genotypes of 11,069 individuals. *Journal of Human Genetics*, epub ahead of print, doi: 10.1038/s10038-020-00847-0.
- Yamada Y. (2019) *History of the Jomon Period*. Kodansha, Tokyo (in Japanese).
- Yaponesian Genome Project Managing Group (2020) Introduction of Yaponesian Genome Project. *iDarwin*, 0: 19–23.
- Zhang W., Meehan J., Su Z., Ng H.W., Shu M., et al. (2014) Whole genome sequencing of 35 individuals provides insights into the genetic architecture of Korean population. *BMC Bioinformatics*, 15: S6.