

A NON-TOXIGENIC BUT MORPHOLOGICALLY AND PHYLOGENETICALLY DISTINCT
NEW SPECIES OF *PSEUDO-NITZSCHIA*, *P. SABIT* SP. NOV. (BACILLARIOPHYCEAE)¹

Sing Tung Teng

Faculty of Resource Science and Technology, Universiti Malaysia Sarawak, Kota Samarahan, Sarawak 94300, Malaysia

Po Teen Lim

Bachok Marine Research Station, Institute of Ocean and Earth Sciences, University of Malaya, Bachok, Kelantan 16310, Malaysia

Hong Chang Lim

Tunku Abdul Rahman University College, Johor Branch Campus, 85000 Segamat, Johor

María Rivera-Vilarelle

Facultad de Ciencias Químicas, Universidad de Colima Km9 Carretera Colima-Coquimatlán, Coquimatlán, Colima, C.P. 28400

Sonia Quijano-Scheggia

Centro Universitario de Investigaciones Oceanológicas, Universidad de Colima, Carretera Manzanillo-Barra de Navidad Km 19.5, Colonia El Naranjo, C.P 28860, Manzanillo, Colima, Mexico

Yoshinobu Takata

The University of Tokyo, Yayoi 1-1-1, Bunkyo-Ku, Tokyo 113-8657, Japan

Michael A. Quilliam

Measurement Science and Standards, Biotoxin Metrology, National Research Council of Canada, 1411 Oxford Street, Halifax, Nova Scotia, Canada B3H 3Z1

Matthias Wolf

Department of Bioinformatics, Biocenter, University of Wuerzburg, Wuerzburg D-97074, Germany

Stephen S. Bates

Fisheries and Oceans Canada, Gulf Fisheries Centre, P.O. Box 5030, Moncton, New Brunswick, Canada E1C 9B6

and *Chui Pin Leaw*²

Bachok Marine Research Station, Institute of Ocean and Earth Sciences, University of Malaya, Bachok, Kelantan 16310, Malaysia

A new species of *Pseudo-nitzschia* (Bacillariophyceae) is described from plankton samples collected from Port Dickson (Malacca Strait, Malaysia) and Manzanillo Bay (Colima, Mexico). The species possesses a distinctive falcate cell valve, from which they form sickle-like colonies in both environmental samples and cultured strains. Detailed observation of frustules under TEM revealed ultrastructure that closely resembles *P. decipiens*, yet the new species differs by the valve shape and greater ranges of striae and poroid densities. The species is readily distinguished from the curve-shaped *P. subcurvata* by the presence of a central interspace. The morphological distinction is further supported by phylogenetic discrimination. We sequenced and

analyzed the nuclear ribosomal RNA genes in the LSU and the second internal transcribed spacer, including its secondary structure, to infer the phylogenetic relationship of the new species with its closest relatives. The results revealed a distinct lineage of the new species, forming a sister cluster with its related species, *P. decipiens* and *P. galaxiae*, but not with *P. subcurvata*. We examined the domoic acid (DA) production of five cultured strains from Malaysia by Liquid chromatography-mass spectrometry (LC-MS), but they showed no detectable DA. Here, we present the taxonomic description of the vegetative cells, document the sexual reproduction, and detail the molecular phylogenetics of *Pseudo-nitzschia sabit* sp. nov.

Key index words: domoic acid; ITS2; Malaysia; Mexico; *Pseudo-nitzschia sabit*; ribosomal RNA genes; secondary structure; sexual reproduction

¹Received 16 October 2014. Accepted 18 May 2015.

²Author for correspondence: e-mail cpleaw@um.edu.my.
Editorial Responsibility: S. Lin (Associate Editor)