

all thermal strategy (Congdon 1989).

Acknowledgments.—We thank S. Allen, C. Burnley, J. Demuth, and G. Smith for assistance in the field, J. Moon for providing logistical support, and H. and I. Lisle for providing a pleasant environment for writing. Turtles were collected under Scientific Collecting Permit No. 675 issued by the Arkansas Game and Fish Commission. This research was approved by the Harding University Animal Care Committee and followed the guidelines in the booklet, "Guidelines for Use of Live Amphibians and Reptiles in Field Research" (1987; American Society of Ichthyologists and Herpetologists, The Herpetologists' League, and the Society for the Study of Amphibians and Reptiles). This research was supported in part by several Faculty Development grants from Harding University.

LITERATURE CITED

- AUTH, D. L. 1975. Behavioral ecology of basking in the yellow-bellied turtle, *Chrysemys scripta scripta* (Schoepff). Bull. Florida State Mus. Biol. Sci. 20:1–45.
- BOYER, D. R. 1965. Ecology of the basking habit in turtles. Ecology 46:99–118.
- BROWN, G. P., R. J. BROOKS, AND J. A. LAYFIELD. 1990. Radiotelemetry of body temperatures of free-ranging snapping turtles (*Chelydra serpentina*) during summer. Canadian J. Zool. 68:1659–1663.
- CONGDON, J. D. 1989. Proximate and evolutionary constraints on energy relations of reptiles. Physiol. Zool. 62:356–373.
- CRAWFORD, K. M., J. R. SPOTILA, AND E. A. STANDORA. 1983. Operative environmental temperatures and basking behavior of the turtle *Pseudemys scripta*. Ecology 64:989–999.
- DI TRANI, C., AND M. A. L. ZUFFI. 1997. Thermoregulation of the European pond turtle, *Emys obicularis*, in central Italy. Chelonian Cons. Biol. 2:428–430.
- ERNST, C. H., R. W. BARBOUR, AND J. E. LOVICH. 1994. Turtles of the United States and Canada. Smithsonian Institution Press, Washington, D.C.
- GRAHAM, T. E., AND A. A. GRAHAM. 1991. *Trionyx spiniferus spiniferus* (eastern spiny softshell) burying behavior. Herpetol. Rev. 22:56–57.
- , AND ———. 1997. Ecology of the eastern spiny softshell, *Apalone spinifer spiniferus*, in the Lamoille River, Vermont. Chelon. Cons. Biol. 2:363–369.
- HUEY, R. B. 1982. Temperature, physiology, and the ecology of reptiles. In C. Gans and F. H. Pough (eds.), *Biology of the Reptilia*, Vol. 12, pp. 25–91. Academic Press, New York.
- HUTCHISON, V. H. 1979. Thermoregulation. In M. Harless and H. Morlock (eds.), *Turtles: Perspectives and Research*, pp. 207–228. John Wiley and Sons, New York.
- , A. VINEGAR, AND R. J. KOSH. 1966. Critical thermal maxima in turtles. Herpetologica 22:32–41.
- KHOSATZKY, L. I. 1981. Some peculiar features of heat exchange exhibited by the softshelled turtle *Trionyx sinensis* (Weigm.). In L. J. Borkin (ed.), *Herpetological Investigations in Siberia and the Far East*, pp. 113–117. Leningrad.
- KING, J. M., G. KUCHLING, AND S. D. BRADSHAW. 1998. Thermal environment, behavior, and body condition of wild *Pseudemys umbrina* (Testudines: Chelidae) during late winter and early spring. Herpetologica 54:103–112.
- LINDEMAN, P. V. 2001. A contrast in the basking habits of the sympatric trionychid turtles *Apalone mutica* and *A. spiniferus*. Herpetol. Nat. Hist. 8:87–89.
- LITZGUS, J. D., AND R. J. BROOKS. 2000. Habitat and temperature selection of *Clemmys guttata* in a northern population. J. Herpetol. 34:178–185.
- MANNING, B., AND G. G. GRIGG. 1997. Basking is not of thermoregulatory significance in the "basking" freshwater turtle *Emydura signata*. Copeia 1997:579–584.
- MOLL, E. O., AND J. M. LEGLER. 1971. The life history of a neotropical slider turtle, *Pseudemys scripta* (Schoepff), in Panama. Bull. Los Angeles Co. Mus. Nat. Hist. 11:1–102.
- NEBEKER, A. V., AND R. B. BURY. 2001. Response of juvenile softshell turtles (*Apalone mutica*) in a thermal gradient. Chelonian Cons. Biol. 4:162–165.
- PLUMMER, M. V. 1977. Activity, habitat and population structure in the turtle, *Trionyx muticus*. Copeia 1977:431–440.
- , AND J. C. BURNLEY. 1997. Behavior, hibernacula, and thermal relations of softshell turtles (*Trionyx spiniferus*) overwintering in a small stream. Chelonian Cons. Biol. 2:489–493.
- , N. E. MILLS, AND S. L. ALLEN. 1997. Activity, habitat, and movement patterns of softshell turtles (*Trionyx spiniferus*) in a small stream. Chelonian Cons. Biol. 2:514–520.
- ROBERTSON, S. L., AND E. N. SMITH. 1982. Evaporative water loss in the spiny soft-shelled turtle *Trionyx spiniferus*. Physiol. Zool. 55:124–129.
- SCHWARZKOPF, L., AND R. J. BROOKS. 1985. Application of operative temperatures to analysis of basking behavior in *Chrysemys picta*. Herpetologica 41:206–212.
- SAJWAJ, T. D., AND J. W. LANG. 2000. Thermal ecology of Blanding's turtle in central Minnesota. Chelonian Cons. Biol. 3:626–636.
- SMITH, E. N., S. L. ROBERTSON, AND S. R. ADAMS. 1981. Thermoregulation of the spiny soft-shelled turtle *Trionyx spiniferus*. Physiol. Zool. 54:74–80.
- SPOTILA, J. R., R. E. FOLEY, AND E. A. STANDORA. 1990. Thermoregulation and climate space of the slider turtle. In J. W. Gibbons (ed.), *Life History and Ecology of the Slider Turtle*, pp. 288–298. Smithsonian Institution Press, Washington, D.C.
- STONE, P. A., AND J. B. IVERSON. 1999. Cutaneous surface area in freshwater turtles. Chelonian Cons. Biol. 3:512–515.
- SYSTAT. 2002. SYSTAT 10.2 Statistics. SYSTAT Software Inc. Richmond, California.
- WILLIAMS, T. A., AND J. L. CHRISTIANSEN. 1981. The niches of two sympatric softshell turtles, *Trionyx muticus* and *Trionyx spiniferus*, in Iowa. J. Herpetol. 15:303–308.

Herpetological Review, 2005, 36(4), 375–382.

© 2005 by Society for the Study of Amphibians and Reptiles

Sources of Larval Identities for Amphibians from Borneo

INDRANEIL DAS

Institute of Biodiversity and Environmental Conservation
Universiti Malaysia Sarawak, 94300, Kota Samarahan
Sarawak, Malaysia
e-mail: idas@ibec.unimas.my

and

ALEXANDER HAAS

Biozentrum Grindel und Zoologisches Museum, Universität Hamburg
Martin-Luther-King-Platz 3, 20146 Hamburg, Germany
e-mail: alexander.haas@uni-hamburg.de

At present, 160 species of amphibians (composed exclusively of Anura and Gymnophiona) are known from Borneo (Das 2005; Das and Haas 2005; Inger and Stuebing 2005). The last summary of knowledge on their larval forms, compiled by Inger (1985), revealed that 45–60% of the fauna then described (63 larval forms), had known larval stages. An additional 14 larval forms could not be assigned to the parent species at the time. Some progress has been made in the two decades that has now elapsed, through the