

**GRAPTEMYS PSEUDOGEOGRAPHICA KOHNII** (Mississippi Map Turtle). USA: TEXAS: COLORADO CO.: Intact and identifiable mummified remains were found amid flood debris, 1.42 km NW of the intersection between FM 949 and Frelsburg Road on FM 949 (29.73949°N, -96.41312°W). 18 June 2006. J. Duvall and M.R.J. Forstner. Verified and accessioned by Toby Hibbits, Curator of Herpetology, Texas Cooperative Wildlife Collection (TCWC 91303). Skin samples were retained in ethanol at Texas State University (MJF 11019). First occurrence west of the Brazos River drainage system and first county record (Dixon 2000. *Amphibians and Reptiles of Texas*. 2<sup>nd</sup> ed. Texas A&M Univ. Press, College Station).

Submitted by **DAVID RODRIGUEZ, JOSEPHINE DUVALL**, and **MICHAEL R.J. FORSTNER**, Texas State University, San Marcos, Texas 78666, USA (e-mail: daverdz5@yahoo.com).

**KINOSTERNON SUBRUBRUM HIPPOCREPIS** (Mississippi Mud Turtle). USA: TEXAS: ARANSAS CO.: 28°14'18.5"N, 67°48'15.3"W. 01 August 2006. Curtis Jones. Verified by Lee Fitzgerald. New county record (Dixon et al. 2000. *Amphibians and Reptiles of Texas*. Texas A&M University Press, College Station. 195 pp. Map 45). Adult found crossing road on the Aransas National Wildlife Refuge. Because of recent rain events there was an abundance of standing water.

Submitted by **CURTIS JONES**, Aransas National Wildlife Refuge, P.O. Box 100 Austwell, Texas 77950, USA.

**LEPIDOCHELYS KEMPII** (Kemp's Ridley Seaturtle). USA: FLORIDA: BREVARD CO.: Canaveral National Seashore. Playalinda Beach (UTM 17 533727 E, 317327 N). 13 June 2006. Florida Museum of Natural History photo voucher (UF 149613). Verified by Steve Johnson. This is the first record of a Kemp's Ridley nesting in Brevard Co. and only the fifth documented individual to nest on the east coast of Florida. This is the second Kemp's Ridley to nest in Canaveral National Seashore, and is 24.8 km S of the first record in 2003 (Fish and Wildlife Institute, Florida Fish and Wildlife Conservation Commission. 19 February 2006. Reported Nesting Activity of Kemp's Ridley, *Lepidochelys kempii* in Florida 1975–2005).

Submitted by **ROBERT H. HEGNA**, Department of Biology, Olivet Nazarene University, Bournbournais, Illinois 60914, USA (e-mail: RHegna@aol.com); and **MATTHEW J. WARREN, CANDACE J. CARTER** (e-mail: Candace\_Carter@nps.gov), and **JOHN C. STINER** (e-mail: John\_Stiner@nps.gov), Canaveral National Seashore, Titusville, Florida 32796, USA.

**MACROCHELYS TEMMINCKII** (Alligator Snapping Turtle). USA: LOUISIANA: MOREHOUSE PARISH: Bayou Bonne Idee near the end of Journey's End Road. 11 April 1997. Joe Evans. Verified by John L. Carr. Natural History Museum, University of Kansas (KU CT 11925, photo voucher, carapace only). First parish record (Dundee and Rossman 1989. *The Amphibians and Reptiles of Louisiana*, Louisiana St. Univ. Press, Baton Rouge. 300 pp.).

Submitted by **GEORGE M. PATTON** and **MARTHA ANN MESSINGER**, 2022 Gemini Drive, Bastrop, Louisiana 71220-3467, USA (e-mail: gpatton@bayou.com).

**PELODISCUS SINENSIS** (Chinese Softshell Turtle). MALAYSIA: SARAWAK: FIRST DIVISION: Sungei Sarawak: Kuching (01°35'N, 110°20'E). 2001. K. A. Jensen and I. Das. USDZ 2.4931. Verified by Kelvin K. P. Lim. First specific locality. This species has been seen for sale on the streets of downtown Kuching and on five occasions found traveling from one canal to another during rainstorms, between 2001–2006. It has been cultured for food at one licensed locality in Kuching for over 10 years. Presence of this soft-shell turtle in the rivers and canals around Kuching indicates that there is an established local population. The species is native to temperate Asia including China, Vietnam, Taiwan, and Hainan, and the Amur River of Russia (Ananjeva et al. 1998. *Amphibians and Reptiles*. Encyclopedia of the Nature of Russia. ABF Press, Moscow. 574 pp.), and introduced to a number of places around tropical Southeast Asia, primarily as escapees from turtle farms (Lim and Das 1999. *Turtles of Borneo and Peninsular Malaysia*. Natural History Publications [Borneo] Sdn. Bhd., Kota Kinabalu. xii + 151 pp.).

Submitted by **KAREN A. JENSEN** (e-mail: kitti\_jensen@yahoo.com), and **INDRANEIL DAS**, Institute of Biodiversity and Environmental Conservation, Universiti Malaysia Sarawak, 94300, Kota Samarahan, Sarawak, Malaysia (e-mail: idas@ibec.unimas.my).

**STERNOTHERUS CARINATUS** (Razor-backed Musk Turtle). USA: TEXAS: TRINITY CO.: Specimen found dead from unknown causes 500 m NE of the Trinity River under an elevated portion of State Highway 19 (30°52'24.4"N, 95°23'10.9"W). 23 September 2006. TCWC 91576. Verified by Toby Hibbits. New county record (Dixon 2000. *Amphibians and Reptiles of Texas*, Second edition, Texas A&M Univ. Press, College Station. 421 pp.). Partially fills in the hiatus in east Texas between Houston and Tyler counties.

Submitted by **KJ LODRIGUE, JR.**, Texas A&M University, Department of Wildlife & Fisheries Sciences, College Station, Texas 77843-2258, USA; and **GRAHAMS S. CRIGLOW**, Brazos Valley Museum of Natural History, 3232 Briarcrest Drive, Bryan, Texas 77802, USA.

**STERNOTHERUS ODORATUS** (Stinkpot). USA: LOUISIANA: MOREHOUSE PARISH: Irvin's Lake (32°57'N, 91°56'W). February 1955. G. M. Patton. Verified by John L. Carr. Field Museum of Natural History (FMNH 246797). First parish record (Dundee and Rossman 1989. *The Amphibians and Reptiles of Louisiana*, Louisiana St. Univ. Press, Baton Rouge. 300 pp.).

Submitted by **GEORGE M. PATTON** and **MARTHA ANN MESSINGER**, 2022 Gemini Drive, Bastrop, Louisiana 71220-3467, USA (e-mail: gpatton@bayou.com).

**TRACHEMYS SCRIPTA ELEGANS** (Red-eared Slider). USA: TEXAS: MADISON CO.: Specimen collected in baited trap ca. 2.7 km S of Madisonville, at the intersection of State Highway 90 and County Road 106 in Caney Creek (30°54'41.7"N, 95°54'38.4"W). 23 September 2006. TCWC 91580. Verified by Toby Hibbits. New county record (Dixon 2000. *Amphibians and Reptiles of Texas*, Second edition, Texas A&M Univ. Press, College Station. 421 pp.). Fills in the hiatus between Grimes and Leon counties.

Submitted by **KJ LODRIGUE, JR.**, Texas A&M University, Department of Wildlife & Fisheries Sciences, College Station,