

# *Herpetological Review*

Volume 40, Number 3 — September 2009





*granulosa* occupying the North Fork Feather River watershed, a distinct watershed from the West Branch Feather River watershed, previously considered the southern extent of the species range in the Sierra Nevada of California (Hayes and Cliff, *op. cit.*). *Taricha granulosa* is found in sympatry with *Taricha torosa* at localities provided herein.

Submitted by **KEVIN D. WISEMAN**, Garcia and Associates (GANDA), 2601 Mission Street, Suite 600, San Francisco, California 94110, USA; Department of Herpetology, California Academy of Sciences, 55 Music Concourse Drive, Golden Gate Park, San Francisco, California 94118, USA; e-mail: kwiseman@garciaandassociates.com.

## ANURA – FROGS

**AGALYCHNIS SPURELLI** (Gliding Treefrog). COSTA RICA: SAN JOSE: Distrito Carara, El Sur de Turrubares, (9.4339°N, 84.34038°W; WGS84), 385 m elev. 19 August 2006. A. Vega, J. M. Robertson, and G. Chaves. Verified by Federico Bolaños. Museum of Zoology, Universidad de Costa Rica, San Jose (UCR 19214). New province record and a 114 km NW range extension from the closest previous record at Palmar Norte, Puntarenas, expanding its distribution along the Pacific versant on the Cordillera de Talamanca (Savage 2002. *The Amphibians and Reptiles of Costa Rica: A Herpetofauna Between Two Continents, Between Two Seas*. Univ. Chicago Press, Chicago, Illinois. xx + 934 pp.). The frog was calling from a tree-lined temporary pool in an agricultural field once covered by lowland rainforest, along with a large breeding congregation of *Agalychnis callidryas*.

Submitted by **ANDRES VEGA**, AMBICOR, 400 E., 75 S., 75 E. de la Municipalidad de Tibas, Tibas, Costa Rica (e-mail: avegak@hotmail.com); **JEANNE M. ROBERTSON**, Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, New York 14853, USA (e-mail: jmr79@cornell.edu); and **GERARDO CHAVES**, Museum of Zoology, University of Costa Rica, San Jose (e-mail: cachi@biologia.ucr.ac.cr).

**ANAXYRUS DEBILIS DEBILIS** (Eastern Green Toad). USA: TEXAS: STEPHENS CO.: N of Eolian (32.68165°N, 99.03378°W; 370 m elev.). 21 March 2009. J. Meik, J. Streicher, and S. Schaack. Verified by C. J. Franklin. University of Texas at Arlington, UTA A-59175. Collected under a rock adjacent to a gravel road in Post Oak savannah. First known county record of the species, which fills a distributional gap in north-central Texas (Dixon 2000. *Amphibians and Reptiles of Texas*. 2<sup>nd</sup> ed. Texas A&M Univ. Press, College Station). Collected under Texas Parks and Wildlife permit number SPR-0707-1387.

Submitted by **JEFFREY W. STREICHER** (e-mail: streicher@uta.edu.), **JESSE M. MEIK**, and **SARAH SCHAACK**, Department of Biology, The University of Texas at Arlington, Arlington, Texas 76019, USA.

**HYLA CINEREA** (Green Treefrog). USA: TENNESSEE: VAN BUREN CO.: Fall Creek Falls State Park Golf Course Pond (35.65°N, 85.3606°W: datum NAD 27). 11 July 2008. Matthew L. Brown and Leigh Ann Brown. Verified by A. Floyd Scott. One adult photographed and catalogued at Austin Peay State University Center for Field Biology (APSU 19959). Only specimen found in park.

Calls have been audible for at least two years at the Fall Creek Falls State Park Golf Course pond. Possible introduction, 62.2 km N of nearest recorded occurrence (Miller et al. 2007. *Herpetol. Rev.* 38:97). First county record (Redmond and Scott 1996. *Atlas of Amphibians in Tennessee*. Misc. Publ. No. 12. The Center for Field Biology, Austin Peay State University, Clarksville, Tennessee. 94 pp. Internet version [http://www.apsu.edu/amatlas] contains links to information regarding Tennessee distribution of amphibians recorded since 1996; accessed 17 January 2009).

Submitted by **MATTHEW L. BROWN**, Fall Creek Falls State Park, Pikeville, Tennessee 37367, USA; e-mail: matthew.brown@state.tn.us.

**GASTROPHRYNE CAROLINENSIS** (Eastern Narrow-mouthed Toad). USA: ARKANSAS: BOONE CO.: 3.2 km SE of Harrison off US 65 (36.122014°N, 93.039479°N; NAD83). 10 June 1995. H. W. Robison. Verified by S. E. Trauth. Arkansas State University Herpetological Museum (ASUMZ 31305). New county record partially filling a distributional hiatus in northern Arkansas between Benton and Marion counties (Trauth et al. 2004. *Amphibians and Reptiles of Arkansas*. Univ. Arkansas Press, Fayetteville. 421 pp.). This frog has now been documented in 69 of 75 (92%) Arkansas counties.

Submitted by **CHRIS T. McALLISTER**, RapidWrite, 102 Brown Street, Hot Springs National Park, Arkansas 71913, USA (e-mail: drcmcallister@aol.com); and **HENRY W. ROBISON**, Department of Biology, Southern Arkansas University, Magnolia, Arkansas 71754, USA (e-mail: hwrobison@suddenlink.net).

**KALOULA PULCHRA** (Painted Bull Frog): MALAYSIA: SARAWAK: MIRI DIVISION: Kuala Lutong, nr. Lutong Town (4.4708333°N, 114.0038889°E). From beach forest, within a patch of *Casuarina* with blackwater pools. 25 October 2007. I. Das. USDZ 1.12153. Verified by Kelvin K. P. Lim. First record for Sarawak. Bornean records from Sinkawang, Kalimantan (Indonesia) by Inger (1966. *Fieldiana Zool.* 52:1–402), Sabah East Malaysia, without locality (Inger and Tan 1996. *Raffles Bull. Zool.* 44:551–574), Crocker Range, Sabah (Das 2006. *Amphibian and Reptile Conservation* 4[1]:3–11), and Bandar Seri Begawan, in Brunei Darussalam (Charles 2008. *Herpetol. Rev.* 39:462–463). Sarawak locality is ca. 115 km W of Brunei record, and choruses of breeding males heard during late afternoons and nights of 25–26 October 2007 indicate species is established locally. Considered introduced by anthropogenic activities in Borneo and is currently spreading west along north coast, and widespread in Southeast Asian mainland (Inger 1999. *In W. E. Duellman [ed.], Patterns of Distribution of Amphibians: A Global Perspective*, pp. 445–482. The John Hopkins University Press, Baltimore and London).

I thank Lee Nyanti, Andrew Alek Tuen, and Matteo Veronesi for supporting field work in Lutong under Project Kuala for Shell Sarawak Berhad and Environmental Resources Management Sdn. Bhd.

Submitted by **INDRANEIL DAS**, Institute of Biodiversity and Environmental Conservation, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia; e-mail: idas@ibec.unimas.my.