



CODEN [USA]: IAJPBB

ISSN: 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

<http://doi.org/10.5281/zenodo.3459188>

Available online at: <http://www.iajps.com>

Research Article

MICROSTRUCTURAL CHANGES IN THE THYROID GLAND OF THE PREWEANING AND WEANING RATS AFTER HETEROTYPIC STRESS

¹Marina Kapitonova, ²Azhar Ahmad, ³Syed Baharom Syed Ahmad Fuad, ⁴Sergey Gupalo,
⁵Sergey Lvovich Kuznetsov, ⁶Sergey Sergeevich Dydykin

¹Full Professor of Anatomy, Faculty of Medicine and Health Sciences, Universiti Malaysia Sarawak (UNIMAS), Kota Samarahan, 94300 Sarawak, Malaysia, e-mail: marinakapitonova@mal.ru, mob:+6017-6243699; ²Science Officer, Faculty of Medicine, Universiti Teknologi MARA, Jalan Hospital, Sungai Buloh 47000, Selangor, Malaysia, e-mail: biotech.azhar@gmail.com; ³MD, MSc (Anatomy), Head of Anatomy Department, Faculty of Medicine, Universiti Teknologi MARA, UiTM Medical and Dental campus, Jalan Hospital, Sungai Buloh 47000, Selangor, Malaysia, e-mail: sbsaf11@gmail.com, ⁴MD, MSc (Pathology), Faculty of Medicine, UNISZA, Kuala Terengganu 21300, Terengganu, Malaysia, e-mail: sgupalo@yandex.ru; ⁵Head of Histology, Cytology and Embryology Department, Sechenov University, 11, Mokhovaya Street, Moscow 125009, Russia, e-mail: vakmedbiol@rambler.ru; ⁶Head of Clinical Anatomy and Operative Surgery Department, Sechenov University, 11, Mokhovaya Street, Moscow 125009, Russia, e-mail: dydykin_ss@mail.ru.

Article Received: July 2019

Accepted: August 2019

Published: September 2019

Abstract:

Chronic stress modulates the activity of various neuroendocrine axes; while activation pattern of the hypothalamo-hypophyseo-adrenal axis (HHAA) is studied in details, the response of the hypothalamo-hypophyseo-thyroid axis (HHTA) is less understood, especially in terms of its response to different types of stressors by the follicular and parafollicular compartments of the thyroid gland. Recent papers presented data on the inhibitory effect of chronic stress on the HHTA, while exercise and low temperature were shown to be able to activate its function. Most studies were done on the adult experimental animals using genetic studies without consideration of the structural changes in the peripheral link of the HHTA in the growing body. Reports on the effect of the chronic variable stressors on the thyroid gland in early life are scarce, though during this period neuroendocrine axes are extremely sensitive to different adverse factors, such as stress, infection, inflammation and environmental changes. The objective of this research is to assess the microscopic changes in the thyroid gland of the preweaning and weaning rats exposed to chronic heterotypic stress compared to the homotypic one. Homo- or heterotypic stressors were chronically applied to the preweaning and weaning rat pups. After the end of the last stress session the animals were euthanized, thyroid gland was sampled, embedded in paraffin, sectioned and stained for thyroglobulin, calcitonin, proliferative cells nuclear antigen (PCNA) and caspase 3. The mucosa of the alimentary tract of the experimental animals was examined and the thymus and the adrenal glands were sampled and weighed to evaluate the depth of stress-induced changes in the body. Immunologically stained slides of the thyroid gland were assessed using Image Pro+ software. Our study showed that chronic stress resulted in the structural and immunohistochemical changes of the thyroid gland in the preweaning and weaning experimental animals which indicate an inhibition of its function in the type of stress-related pattern. Both homo- and heterotypic stressors caused microscopic alterations in the thyroid gland, the extent of which depended both on the initial age of the experimental animal and the type of the stressor applied. The number and the size of the thyroglobulin-positive cells significantly decreased, and the volume density of the apoptotic cells significantly increased in the heterotypically stressed rat pups of both age subgroups with higher level of significance in the preweaning age subgroup. The number of calcitoninocytes was significantly increased in the weaning rat pups exposed to the heterotypic stress. The number of PCNA-positive cells significantly decreased only in the heterotypic stress group of both ages with higher level of significance in the preweaning age subgroup. A positive correlation was found between the volume density of the thyroglobulin-positive cells and the severity of the accidental thymic involution. Thus, our research demonstrated that weaning period is very sensitive to chronic stress for the thyroid gland of the experimental animals which by this age becomes mature enough to differentially respond to the various types of stress (homotypic versus heterotypic) both by the thyroid follicular and parafollicular compartments, and that thymus plays an important role in the functional capacity of the thyroid gland during stress in early life.

Key words: Thyroid Gland, Chronic Stress, Weaning Period, Immunohistochemistry, Image Analysis

9. Sotelo-Rivera I, Jaimes-Hoy L, Cote-Vélez A, Espinoza-Ayala C, Charli JL, Joseph-Bravo P. An acute injection of corticosterone increases thyrotrophin-releasing hormone expression in the paraventricular nucleus of the hypothalamus but interferes with the rapid hypothalamus pituitary thyroid axis response to cold in male rats. *J Neuroendocrinol*, 2014;26:861-9.
10. Gutiérrez-Mariscal M, de Gortari P, López-Rubalcava C, Martínez A, Joseph-Bravo P. Analysis of the anxiolytic-like effect of TRH and the response of amygdalar TRHergic neurons in anxiety. *Psychoneuroendocrinology*, 2008;33:198-213.
11. Aguilar-Valles A, Sánchez E, de Gortari P, García-Vazquez AI, Ramírez-Amaya V, Bermúdez-Rattoni F, Joseph-Bravo P. The expression of TRH, its receptors and degrading enzyme is differentially modulated in the rat limbic system during training in the Morris water maze. *Neurochem Int*, 2007;50:404-17.
12. Armario A, Marti O, Gavaldà A, Giralt M, Jolin T. Effects of chronic immobilization stress on GH and TSH secretion in the rat: response to hypothalamic regulatory factors. *Psychoneuroendocrinology*, 1993;18:405-13.
13. Cizza G, Brady LS, Esclapes ME, Blackman MR, Gold PW, Chrousos GP. Age and gender influence basal and stress-modulated hypothalamic-pituitary-thyroidal function in Fischer 344/N rats. *Neuroendocrinology*, 1996;64:440-8.
14. Baumgartner A. Thyroxine and the treatment of affective disorders: An overview of the results of basic and clinical research. *Int J Neuropsychopharmacol*, 2000;3:149-65.
15. Bianco AC, Maia AL, da Silva WS, Christoffolete MA. Adaptive activation of thyroid hormone and energy expenditure. *Biosci Rep*, 2005;25:191-208.
16. Egri P, Fekete C, Dénes Á, Reglódi D, Hashimoto H, Fülöp BD, Gereben B. Pituitary Adenylate Cyclase-Activating Polypeptide (PACAP) Regulates the Hypothalamo-Pituitary-Thyroid (HPT) Axis via Type 2 Deiodinase in Male Mice. *Endocrinology*, 2016;157:2356-66.
17. Mohácsik P, Füzesi T, Doleschall M, Szilvási-Szabó A, Vancamp P, Hadadi É, Darras VM, Fekete C, Gereben B. Increased Thyroid Hormone Activation Accompanies the Formation of Thyroid Hormone-Dependent Negative Feedback in Developing Chicken Hypothalamus. *Endocrinology*. 2016 Mar;157:1211-21.
18. Vieira, JO, Duarte, JO, Costa-Ferreira W, Morais-Silva G, Marin MT, Crestani CC. Sex differences in cardiovascular, neuroendocrine and behavioral changes evoked by chronic stressors in rats. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 2018; 81:426-37.
19. Umeoka EHL, Robinson EJ, Turimella SL, van Campen JS, Motta-Teixeira RLC, Sarabdjitsingh A, Garcia-Cairasco N, Braun K, de Graan PN, Joëls M. Hyperthermia-induced seizures followed by repetitive stress are associated with age-dependent changes in specific aspects of the mouse stress system. *Journal of Neuroendocrinology*, 2019; e12697.
20. Klein JR. The immune system as a regulator of thyroid hormone activity. *Exp Biol Med (Maywood)*, 2006;231:229-36.
21. Choudhury BK, Shi XZ, Sarna SK. Norepinephrine mediates the transcriptional effects of heterotypic chronic stress on colonic motor function. *Am J Physiol Gastrointest Liver Physiol*, 2009;296:1238-47.
22. Khmel'nitskiĭ OK, Khmel'nitskaia NM, Tararak TIa, Vasil'eva NA, Balykin MV. Impact of intermittent hypobaric hypoxia on the thyroid morphofunctional state in experimental hyperthyroidism. *Arkh Patol*, 2006;68:31-3.
23. Tsujio M, Watahiki Y, Yoshioka K, Mutoh K. Morphology of thyroid follicular cells of methimazole-treated rats. *Anat Histol Embryol*, 2007;36:290-4.
24. Zbucki RL, Winnicka MM, Sawicki B, Szynaka B, Andrzejewska A, Puchalski Z. Alteration of parafollicular (C) cells activity in the experimental model of hypothyroidism in rats. *Folia Histochem Cytobiol*, 2007;45:115-21.