

**Running head: TRAUMA EXPOSURE BEHAVIOR AND PHYSIOLOGICAL
MEASURE**

**Lifetime Trauma Exposure, and PTSD Symptoms in relation to Health-related
Behaviors and Physiological Measures among Malaysian Adolescents**

Yoke Yong Chen, PhD and Siti Raudzah Ghazali, PhD

Universiti Malaysia Sarawak

Author Note

Yoke Yong Chen, Siti Raudzah Ghazali, Faculty of Medicine and Health Science,

Universiti Malaysia Sarawak

Correspondence concerning this article should be addressed to Chen Yoke Yong,
Department of Psychological Medicine, Faculty of Medicine and Health Science, Universiti
Malaysia Sarawak, Sarawak, Kota Samarahan 94300 Sarawak, MALAYSIA. Email:
yongcyy@gmail.com

This research was funded by The Ministry of Higher Education Malaysia, RACE/e
(1)/888/2012 (06).

This article has not been published elsewhere and has not also been submitted simultaneously
for publication elsewhere and there is no conflict of interest.

TRAUMA EXPOSURE BEHAVIOR AND PHYSIOLOGICAL MEASURE 1**Abstract**

Lifetime trauma exposure may result in serious consequences for mental health. It can lead to Posttraumatic Stress Disorder (PTSD) diagnosis, and when traumatic experience occurs among adolescents the consequences may affect their mental and physical health later in their adulthood. This is a cross-sectional research design aiming to determine the relationship between lifetime trauma experience with health-related behaviors and physiological measures among Malaysian adolescents. Six-hundred and six adolescents with mean age of 16.9 years ($SD=1.28$) responded to the survey questionnaires. Their height, weight, blood pressure and heart rate were measured. Adolescents with traumatic experiences were significantly more likely to engage in unhealthy behaviors such as drinking alcohol and smoking. Their physiological readings (i.e., heart rate) were significantly higher than those without traumatic experience. The significant relationship between trauma exposure, unhealthy behavior and physical health are discussed in this paper.

Keywords: *Lifetime trauma, health-related behaviors, physiological measures, adolescents, PTSD*

Lifetime Trauma Exposure and PTSD Symptoms in relation to Health-related Behavior and Physiological Measures among Malaysian Adolescents

Childhood traumatic experiences may potentially lead to long term psychological problems (Hughes et al., 2017; Scott et al., 2013). These traumatic experiences contribute to poorer life-course health outcomes (Bellis, Lowey, Leckenby, Hughes, & Harrison, 2014; May-Ling et al., 2015). This may due to the enduring changes in human nervous and endocrine systems and imbalanced allostatic systems, which can lead to progressive physiological damage (Danese & McEwen, 2012; Tulloch, Greenman, & Tassé, 2014). Individuals who experience traumatic events are prone to adopting unhealthy behaviors and are more susceptible to non-communicable diseases. Unfortunately, studies on the increased risk for poor physical health outcomes due to trauma exposure, and the development and manifestation of PTSD have only been investigated and reported in the last few decades (Coughlin, 2011).

The relationship between trauma exposure and/or posttraumatic stress disorder (PTSD) and poor physical outcomes such as cardiovascular disease (CVD) can be explained by sympathetic overactivation theory (Danese & McEwen, 2012). Overactivation of sympathetic responses results in an elevated heart rate which in turn is significantly associated with overweight and obesity (Danese & McEwen, 2012; Gandubert et al., 2016). Overweight and obesity serve as mediators to increase sodium retention in blood vessels and subsequently increase blood pressure which have an adverse effect on physical health (da Silva, do Carmo, Dubinion & Hall, 2009).

A large longitudinal study of 38,352 participants found that PTSD was significantly associated with weight gain and the development of obesity (LeardMann et al., 2015). Overweight individuals were significantly associated with elevated C-