

**NOTE ON THE DISTRIBUTION OF
PTEROSPERMUM PECTENIFORME KOSTERM.
(DOMBEYOIDEAE : MALVACEAE S .L.) IN
PENINSULAR MALAYSIA**

AHMAD FITRI, Z., *¹ NIZAM, M. S., ^{1,2} SAIYIDAH, R. B., ³
ROHAIZA, D., ⁴ FAEZAH, P., ⁵ NURUL-SYAHIRAH, M., ¹
NORAZLINDA, M., ¹ RUZI, A. R., ¹ NUR ‘AQILAH, M. B., ¹ NIK
HAZLAN, N. H., ⁶ NIK NORAFIDA, N. A.¹ AND A. LATIFF¹

¹Department of Biological Sciences and Biotechnology, Faculty of Science and Technology, Universiti Kebangsaan Malaysia 43600, Bangi, Selangor

²Institute of Climate Change, Universiti Kebangsaan Malaysia, 43600 UKM Bangi, Selangor, Malaysia

³UiTM Cawangan Sarawak Kampus Mukah, Km 7.5 Jalan Mukah-Oya 96400 Mukah Sarawak, Oya, 96300 Mukah, Sarawak.

⁴Centre for Pre-University Studies, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia.

⁵Faculty of Applied Sciences, Universiti Teknologi MARA, 40450 Shah Alam, Selangor, Malaysia

⁶Centre for Wood Industry Studies, Faculty of Applied Science, Universiti Teknologi MARA Cawangan Pahang, 26400 Bandar Tun Abdul Razak, Jengka, Pahang.

*Corresponding author:

Email: ahmadfitri@ukm.edu.my

Abstract : The rare species *Pterospermum pecteniforme* previously is only known from two herbarium collections in the late 1960 and early 1970 in Perlis and Kedah (Langkawi Islands; Pulau Timun), respectively. In this paper, we have added two additional localities from two islands of Langkawi Archipelago namely, Pulau Dayang Bunting and Pulau Singa Besar. This species is locally quite common on both islands. All fertile collections of this species are only from the fruits while the flower is still unknown.

Key words: *Pterospermum pecteniforme*, Dombeyoideae, Malvaceae s. l.) rare, distribution

INTRODUCTION

Pterospermum was described by Johann Christian Daniel von Schreber, a German professor and naturalist in *Genera Plantarum* in 1791. The ethymology of *Pterospermum* is seeded wing (*ptero*; Greek, = wing; *spermum*; Greek, = seed). This genus comprises of ca. 20-30 species (Wilkie & Berhaman, 2011; Wilkie, 2013) with its distribution from India to South China through Southeast Asia to Maluku (Kochummen, 1973; Wilkie, 2013). In Peninsular Malaysia, six species are recorded