

# The Effect of Vaccination Coverage and Climate on Japanese Encephalitis in Sarawak, Malaysia

Daniel E. Impoinvil<sup>1,2\*</sup>, Mong How Ooi<sup>3</sup>, Peter J. Diggle<sup>1,4</sup>, Cyril Caminade<sup>5</sup>, Mary Jane Cardosa<sup>6</sup>, Andrew P. Morse<sup>5</sup>, Matthew Baylis<sup>1,7</sup>, Tom Solomon<sup>2,7,8</sup>

**1** Liverpool University Climate and Infectious Disease of Animals Group, Department of Epidemiology and Population Health, Institute of Infection and Global Health, University of Liverpool, Neston, Cheshire, United Kingdom, **2** Brain Infections Group, Department of Clinical Infection, Microbiology and Immunology, Institute of Infection and Global Health, University of Liverpool, Merseyside, Liverpool, United Kingdom, **3** Department of Paediatrics, Sibul Hospital, Sibul, Sarawak, Malaysia, **4** Lancaster Medical School, Faculty of Health and Medicine, Lancaster University, Lancaster, United Kingdom, **5** Department of Geography and Planning, School of Environmental Sciences, University of Liverpool, Merseyside, Liverpool, United Kingdom, **6** Institute of Health and Community Medicine, Universiti Malaysia Sarawak, Kota Samarahan, Sarawak, Malaysia, **7** The National Consortium for Zoonosis Research, Neston, Cheshire, United Kingdom, **8** Walton Centre NHS Foundation Trust, Fazakerley, Liverpool, Merseyside, United Kingdom

## Abstract

**Background:** Japanese encephalitis (JE) is the leading cause of viral encephalitis across Asia with approximately 70,000 cases a year and 10,000 to 15,000 deaths. Because JE incidence varies widely over time, partly due to inter-annual climate variability effects on mosquito vector abundance, it becomes more complex to assess the effects of a vaccination programme since more or less climatically favourable years could also contribute to a change in incidence post-vaccination. Therefore, the objective of this study was to quantify vaccination effect on confirmed Japanese encephalitis (JE) cases in Sarawak, Malaysia after controlling for climate variability to better understand temporal dynamics of JE virus transmission and control.

**Methodology/principal findings:** Monthly data on serologically confirmed JE cases were acquired from Sibul Hospital in Sarawak from 1997 to 2006. JE vaccine coverage (non-vaccine years vs. vaccine years) and meteorological predictor variables, including temperature, rainfall and the Southern Oscillation index (SOI) were tested for their association with JE cases using Poisson time series analysis and controlling for seasonality and long-term trend. Over the 10-years surveillance period, 133 confirmed JE cases were identified. There was an estimated 61% reduction in JE risk after the introduction of vaccination, when no account is taken of the effects of climate. This reduction is only approximately 45% when the effects of inter-annual variability in climate are controlled for in the model. The Poisson model indicated that rainfall (lag 1-month), minimum temperature (lag 6-months) and SOI (lag 6-months) were positively associated with JE cases.

**Conclusions/significance:** This study provides the first improved estimate of JE reduction through vaccination by taking account of climate inter-annual variability. Our analysis confirms that vaccination has substantially reduced JE risk in Sarawak but this benefit may be overestimated if climate effects are ignored.

**Citation:** Impoinvil DE, Ooi MH, Diggle PJ, Caminade C, Cardosa MJ, et al. (2013) The Effect of Vaccination Coverage and Climate on Japanese Encephalitis in Sarawak, Malaysia. PLoS Negl Trop Dis 7(8): e2334. doi:10.1371/journal.pntd.0002334

**Editor:** David Joseph Diemert, The George Washington University Medical Center, United States of America

**Received:** March 6, 2013; **Accepted:** June 11, 2013; **Published:** August 8, 2013

**Copyright:** © 2013 Impoinvil et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work's original data collection was supported by the Ministry of Science, Technology and Innovation (NBD06-05-01-T001)(URL: www.mosti.gov.my/), operational funds of Sarawak Health Department (URL: http://jknsarawak.moh.gov.my/en/), the Walton Centre for Neurology and Neurosurgery Research Fund (URL: http://www.thewaltoncentre.nhs.uk/), Program for Appropriate Technology in Health (PATH; Seattle,WA)(www.path.org/), Wellcome Trust Clinical Training Fellowship (awarded to MHO) (www.wellcome.ac.uk/), and a United Kingdom Medical Research Council Senior Clinical Fellowship (awarded to TS) (www.mrc.ac.uk/). The analysis of the study was supported by the Leverhulme Trust Research Leadership Award: "Predicting the effects of climate change on infectious diseases of animals" (awarded to MB) (URL: www.leverhulme.ac.uk/). The funders had no role in study design, data collection and analysis, or preparation of the manuscript. The Ministry of Health approved the publication of this manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: xda6@cdc.gov

† Current address: Centers for Disease Control and Prevention, Division of Parasitic Diseases and Malaria/Entomology Branch, Atlanta, Georgia, United States of America.

‡ These authors contributed equally to this work.

## Introduction

Japanese encephalitis (JE) is a mosquito-borne disease which is the leading cause of viral encephalitis across Asia with approximately 70,000 cases a year and 10,000 to 15,000 deaths [1,2]. There are two transmission patterns of JEV in three climatic zones [3,4]: in temperate climate zones such as Japan, Korea and

mainland China and Taiwan, and sub-tropical climate zones such as Nepal, and northerly areas of India, Thailand and Vietnam, JEV exhibits an epidemic or outbreak transmission pattern, characterized by a well-defined seasonal peak. In tropical climate zones such as Malaysia, Indonesia, southerly areas of Vietnam and Thailand, JEV exhibits an endemic transmission characterized by sporadic human cases throughout the year with seasonal peaks.